

National Institute for Health and Care Excellence

**NICE clinical guideline on suspected cancer
Document cover sheet**

Date	Version number	Editor	Action

Suspected cancer: recognition and management of suspected cancer in children, young people and adults

NICE guideline

Draft for consultation, November 2014

If you wish to comment on this version of the guideline, please be aware that all the supporting information and evidence for the 2015 recommendations is contained in the full version of the 2015 guideline. Evidence for the 2005 recommendations is in the full version of the 2005 guideline.'

Contents

Introduction	3
Patient-centred care.....	4
Strength of recommendations	5
Update information.....	7
Recommendations organised by cancer symptom and findings of primary care investigations	8
Abdominal symptoms	9
Bleeding	19
Gynaecological symptoms.....	24
Lumps or masses	25
Neurological symptoms	30
Pain	30
Respiratory symptoms.....	31
Skeletal symptoms	38
Skin or surface symptoms	40
Urological symptoms	42
Non-specific features of cancer	44
Primary care investigations	61
Recommendations organised by site of cancer	68
Recommendations on patient support, safety netting and the diagnostic process	89
Research recommendations	94
Other information	95
The Guideline Development Group, National Collaborating Centre and NICE project team	99
Appendix A: Recommendations from NICE clinical guideline 27 (2005) that have been deleted or changed	103

Introduction

Cancer is an important subject, both in terms of the number of people affected and the impacts on those people and the people close to them. Around one third of a million new cancers are diagnosed annually in the UK, across over 200 different cancer types. Each of these cancer types has different presenting features, though there may be overlap. Approximately one third of the population will develop a cancer in their lifetime. There is considerable variation in referral and testing for possible cancer, which cannot be fully explained by variation in the population.

The identification of people with possible cancer is a primary care task, as the large majority of people first present to a primary care clinician. Therefore, evidence from primary care should inform the identification process. Evidence from secondary care can only study those in whom cancer is already suspected, but cannot tell us when we should suspect cancer. Therefore this guideline is based on evidence from primary care.

This guideline covers people of all ages, including children and young adults. It covers the selection of people for referral or investigation in primary care. The guideline also aims to help people understand what to expect if they have symptoms that may suggest cancer. In addition it aims to help those in secondary care to understanding what services should be provided for people with suspected cancer. Finally, these recommendations are recommendations, not requirements, and they are not intended to override clinical judgement.

The recommendations in this guideline have been organised into three separate sections to help clinicians find the relevant information easily. For those wanting to find recommendations on specific symptoms and primary care investigations, the recommendations are in a section organised by symptoms and investigation findings. The same recommendations are also organised in the following section by cancer site. Finally there is a section covering patient support, safety netting and the diagnostic process.

Patient-centred care

This guideline offers best practice advice on the care of people with suspected cancer.

Patients and healthcare professionals have rights and responsibilities as set out in the [NHS Constitution for England](#) – all NICE guidance is written to reflect these. Treatment and care should take into account individual needs and preferences. Patients should have the opportunity to make informed decisions about their care and treatment, in partnership with their healthcare professionals. If the patient is under 16, their family or carers should also be given information and support to help the child or young person to make decisions about their treatment. Healthcare professionals should follow the [Department of Health's advice on consent](#). If someone does not have capacity to make decisions, healthcare professionals should follow the [code of practice that accompanies the Mental Capacity Act](#) and the supplementary [code of practice on deprivation of liberty safeguards](#).

NICE has produced guidance on the components of good patient experience in adult NHS services. All healthcare professionals should follow the recommendations in [Patient experience in adult NHS services](#).

If a young person is moving between paediatric and adult services, care should be planned and managed according to the best practice guidance described in the Department of Health's [Transition: getting it right for young people](#).

Adult and paediatric healthcare teams should work jointly to provide assessment and services to young people with suspected cancer. Diagnosis and management should be reviewed throughout the transition process, and there should be clarity about who is the lead clinician to ensure continuity of care.

Strength of recommendations

Some recommendations can be made with more certainty than others. The Guideline Development Group makes a recommendation based on the trade-off between the benefits and harms of an intervention, taking into account the quality of the underpinning evidence. For some interventions, the Guideline Development Group is confident that, given the information it has looked at, most patients would choose the intervention. The wording used in the recommendations in this guideline denotes the certainty with which the recommendation is made (the strength of the recommendation).

For all recommendations, NICE expects that there is discussion with the patient about the risks and benefits of the interventions, and their values and preferences. This discussion aims to help them to reach a fully informed decision (see also 'Patient-centred care').

Interventions that must (or must not) be used

We usually use 'must' or 'must not' only if there is a legal duty to apply the recommendation. Occasionally we use 'must' (or 'must not') if the consequences of not following the recommendation could be extremely serious or potentially life threatening.

Interventions that should (or should not) be used – a 'strong' recommendation

We use 'offer' (and similar words such as 'refer' or 'advise') when we are confident that, for the vast majority of patients, an intervention will do more good than harm, and be cost effective. We use similar forms of words (for example, 'Do not offer...') when we are confident that an intervention will not be of benefit for most patients.

Interventions that could be used

We use 'consider' when we are confident that an intervention will do more good than harm for most patients, and be cost effective, but other options may be similarly cost effective. The choice of intervention, and whether or not to have the intervention at all, is more likely to depend on the patient's values

and preferences than for a strong recommendation, and so the healthcare professional should spend more time considering and discussing the options with the patient.

Recommendation wording in guideline updates

NICE began using this approach to denote the strength of recommendations in guidelines that started development after publication of the 2009 version of 'The guidelines manual' (January 2009). This does not apply to any recommendations shaded in grey and ending [2005] (see 'Update information' box below for details about how recommendations are labelled). In particular, for recommendations labelled [2005], the word 'consider' may not necessarily be used to denote the strength of the recommendation.

Update information

This guidance is an update of NICE guideline CG27 (published June 2005) and will replace it.

New recommendations have been added for recognition, management and referral for suspected cancer in primary care.

You are invited to comment on the new and updated recommendations in this guideline. These are marked as:

- **[new 2015]** if the evidence has been reviewed and the recommendation has been added or updated
- **[2015]** if the evidence has been reviewed but no change has been made to the recommended action.

You are also invited to comment on recommendations that NICE proposes to delete from the 2005 guideline, because either the evidence has been reviewed and the recommendations have been updated, or NICE has updated other relevant guidance and has replaced the original recommendations.

Appendix A sets out these recommendations and includes details of replacement recommendations. Where there is no replacement recommendation, an explanation for the proposed deletion is given.

Where recommendations are shaded in grey and end **[2005]**, the evidence has not been reviewed since the original guideline. We will not be able to accept comments on these recommendations.

The original NICE guideline and supporting documents are available [here](#).

1 Recommendations organised by cancer symptom and findings of primary care investigations

The following guidance is based on the best available evidence. The full guideline [hyperlink to be added for final publication] gives details of the methods and the evidence used to develop the guidance.

This section gives details of the recommendations organised by symptoms and findings from primary care investigations. The following categories have been used:

- [Abdominal symptoms](#)
- [Bleeding](#)
- [Gynaecological symptoms](#)
- [Lumps or masses](#)
- [Neurological symptoms](#)
- [Pain](#)
- [Respiratory symptoms](#)
- [Skeletal symptoms](#)
- [Skin or surface symptoms](#)
- [Urological symptoms](#)
- [Non-specific features of cancer](#)
- [Primary care investigations](#)

Abdominal symptoms

See also [Bleeding](#) for recommendations on rectal bleeding.

Abdominal distension

Symptom and specific features	Possible cancer	Recommendation
Abdominal distension – persistent (women)	Ovarian	<p>Carry out tests in primary care if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month:</p> <ul style="list-style-type: none"> • persistent abdominal distension (women often refer to this as 'bloating') • feeling full (early satiety) and/or loss of appetite • pelvic or abdominal pain • increased urinary urgency and/or frequency. [1.4.2] [R57] <p>Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61]</p> <p>See also primary care investigations</p>

Abdominal or pelvic mass or enlarged abdominal organ

Symptom and specific features	Possible cancer	Recommendation
Hepatosplenomegaly – unexplained (children and young people)	Leukaemia	Refer children and young people for immediate specialist assessment for leukaemia if they have unexplained petechiae or hepatosplenomegaly. [new 2015] [1.8.3] [R107]
Palpable abdominal mass or unexplained enlarged abdominal organ (children)	Neuroblastoma	Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. [new 2015] [1.10.1] [R117]
Palpable abdominal mass or unexplained enlarged abdominal organ (children)	Wilm's tumour	Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. [new 2015] [1.10.3] [R119]
Pelvic or abdominal mass (women)	Ovarian	Refer the woman urgently ¹ if physical examination identifies ascites and/or a pelvic or abdominal mass (which is not obviously uterine fibroids). [1.4.1] [R56]
Splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]
Upper abdominal mass consistent with stomach cancer	Stomach	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with an upper abdominal mass consistent with stomach cancer. [new 2015] [1.2.6] [R36]
Lower gastrointestinal tract symptoms	Colorectal	Offer a digital rectal examination to people In patients with unexplained symptoms related to the lower gastrointestinal tract. [new 2015] [1.3.9] [Ra3]

¹ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

Hepatosplenomegaly	Leukaemia	Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms: <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]
Upper abdominal mass consistent with an enlarged gall bladder	Gall Bladder	Consider urgent direct access ultrasound (within 2 weeks) to assess for gall bladder cancer in people with an upper abdominal mass consistent with an enlarged gall bladder. [new 2015] [1.2.12] [R42]
Upper abdominal mass consistent with an enlarged liver	Liver	Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for liver cancer in people with an upper abdominal mass consistent with an enlarged liver. [new 2015] [1.2.13] [R43]
Rectal or abdominal mass	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people with a rectal or abdominal mass. [new 2015] [1.3.4] [R47]

Abdominal or pelvic pain

Symptom and specific features	Possible cancer	Recommendation
Abdominal pain without rectal bleeding	Colorectal	Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who: <ul style="list-style-type: none"> • have abdominal pain or • have weight loss or • are aged under 60 years and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.6] [R49]

		See also primary care investigations Offer a digital rectal examination to people with unexplained symptoms related to the lower gastrointestinal tract. [new 2015] [1.3.9] [Ra3]
Lower gastrointestinal tract symptoms	Colorectal	Offer a digital rectal examination to people with unexplained symptoms related to the lower gastrointestinal tract. [new 2015] [1.3.9] [Ra3]
Pelvic or abdominal pain (women)	Ovarian	Carry out tests in primary care if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month: <ul style="list-style-type: none"> • persistent abdominal distension (women often refer to this as 'bloating') • feeling full (early satiety) and/or loss of appetite • pelvic or abdominal pain • increased urinary urgency and/or frequency. [1.4.2] [R57] Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61] See also primary care investigations
Irritable bowel syndrome symptoms (women 50 and over)	Ovarian	Carry out appropriate tests for ovarian cancer in any woman of 50 or over who has experienced symptoms within the last 12 months that suggest irritable bowel syndrome (IBS) ² , because IBS rarely presents for the first time in women of this age. [1.4.5] [R60] Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61] See also primary care investigations
Abdominal pain with weight loss (over 40)	Colorectal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 40 with unexplained weight loss and abdominal pain. [new 2015] [1.3.5] [R48]
Upper abdominal pain with weight loss (55 and over)	Oesophageal	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people: <ul style="list-style-type: none"> • aged 45 and over with dysphagia

² See the NICE guideline on [irritable bowel syndrome in adults](#).

		<p>or</p> <ul style="list-style-type: none"> aged 55 and over with weight loss and upper abdominal pain, or reflux or dyspepsia. [new 2015] [1.2.1] [R29]
Upper abdominal pain with weight loss and nausea or vomiting (40 and over) Upper abdominal pain with weight loss (55 and over)	Stomach	<p>Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who:</p> <ul style="list-style-type: none"> are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or are aged 55 and over with upper abdominal pain, reflux or dyspepsia. [new 2015] [1.2.8] [R38]
Abdominal pain with weight loss (under 55)	Stomach	<p>Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who:</p> <ul style="list-style-type: none"> also have appetite loss or are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more or are aged 55 and over with nausea or vomiting. [new 2015] [1.2.9] [R39]
Abdominal pain with weight loss	Pancreatic	<p>Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms:</p> <ul style="list-style-type: none"> diarrhoea back pain abdominal pain nausea or vomiting constipation new-onset diabetes. [new 2015] [1.2.5] [R33]
Abdominal pain with raised platelet count (55 and over)	Oesophageal	<p>Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people aged 55 or over with:</p> <ul style="list-style-type: none"> weight loss and nausea or vomiting or reflux and nausea or vomiting or upper abdominal pain and raised platelet count. [new 2015] [1.2.3]

		[R31]
Abdominal pain with raised platelet count (55 and over)	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with upper abdominal pain and raised platelet counts. [new 2015] [1.2.11] [R41]
Abdominal pain with rectal bleeding (under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]

Change in bowel habit

Symptom and specific features	Possible cancer	Recommendation
Change in bowel habit (over 60)	Colorectal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained changes in their bowel habit. [new 2015] [1.3.3] [R46]
Change in bowel habit without rectal bleeding	Colorectal	Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who: <ul style="list-style-type: none"> • have abdominal pain or • have weight loss or • are aged under 60 years and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.6] [R49] See also primary care investigations Offer a digital rectal examination to people with unexplained symptoms related to the lower gastrointestinal tract. [new 2015] [1.3.9] [Ra3]
Irritable bowel	Ovarian	Carry out appropriate tests for ovarian

syndrome symptoms (women 50 or over)		cancer in any woman of 50 or over who has experienced symptoms within the last 12 months that suggest irritable bowel syndrome (IBS) ³ , because IBS rarely presents for the first time in women of this age. [1.4.5] [R60] Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61] See also primary care investigations
Change in bowel habit	Ovarian	Consider carrying out tests in primary care if a woman reports unexplained weight loss, fatigue or changes in bowel habit. [1.4.3] [R58] Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61] See also primary care investigations
Change in bowel habit with weight loss (60 and over)	Pancreatic	Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms: <ul style="list-style-type: none"> • diarrhoea • back pain • abdominal pain • nausea or vomiting • constipation • new-onset diabetes. [new 2015] [1.2.5] [R33]
Change in bowel habit with rectal bleeding (under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]

Dyspepsia

³ See the NICE guideline on [irritable bowel syndrome in adults](#).

Symptom and specific features	Possible cancer	Recommendation
Dyspepsia with weight loss (55 and over)	Oesophageal	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people: <ul style="list-style-type: none"> aged 45 and over with dysphagia or aged 55 and over with weight loss and upper abdominal pain, or reflux or dyspepsia. [new 2015] [1.2.1] [R29]
Dyspepsia with weight loss (55 and over)	Stomach	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or are aged 55 and over with upper abdominal pain, reflux or dyspepsia. [new 2015] [1.2.8] [R38]
Dyspepsia with weight loss (under 55)	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> also have appetite loss or are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more or are aged 55 and over with nausea or vomiting. [new 2015] [1.2.9] [R39]

Dysphagia

Symptom and specific features	Possible cancer	Recommendation
Dysphagia (45 and over)	Oesophageal	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people: <ul style="list-style-type: none"> aged 45 and over with dysphagia or aged 55 and over with weight loss and upper abdominal pain, or reflux or dyspepsia. [new 2015] [1.2.1] [R29]
Dysphagia	Stomach	Offer urgent direct access upper gastrointestinal endoscopy (within

		2 weeks) to assess for stomach cancer in people with dysphagia. [new 2015] [1.2.7] [R37]
--	--	---

Abdominal examination findings

Symptom and specific features	Possible cancer	Recommendation
Ascites and/or a pelvic or abdominal mass	Ovarian	Refer the woman urgently ⁴ if physical examination identifies ascites and/or a pelvic or abdominal mass (which is not obviously uterine fibroids. [1.4.1] [R56]

Rectal examination findings

Symptom and specific features	Possible cancer	Recommendation
Prostate findings	Prostate	Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for prostate cancer if their prostate feels malignant on digital rectal examination. [new 2015] [1.5.1] [R72]
Anal mass or anal ulceration	Anal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for anal cancer in people with an unexplained anal mass or unexplained anal ulceration. [new 2015] [1.3.10] [R52]
Rectal mass	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people with a rectal or abdominal mass. [new 2015] [1.3.4] [R47]

Nausea or vomiting

Symptom and specific features	Possible cancer	Recommendation
Nausea or vomiting with upper abdominal pain and weight loss (40 and over)	Stomach	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or are aged 55 and over with upper abdominal pain, reflux or

⁴ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

		dyspepsia. [new 2015] [1.2.8] [R38]
Nausea or vomiting with weight loss (55 and over)	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> • also have appetite loss or • are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more or • are aged 55 and over with nausea or vomiting. [new 2015] [1.2.9] [R39]
Nausea or vomiting with weight loss	Pancreatic	Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms: <ul style="list-style-type: none"> • diarrhoea • back pain • abdominal pain • nausea or vomiting • constipation • new-onset diabetes. [new 2015] [1.2.5] [R33]
Nausea or vomiting with reflux	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with reflux and nausea or vomiting. [new 2015] [1.2.10] [R40]
Nausea or vomiting with reflux	Oesophageal	Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people aged 55 or over with: <ul style="list-style-type: none"> • weight loss and nausea or vomiting or • reflux and nausea or vomiting or • upper abdominal pain and raised platelet count. [new 2015] [1.2.3] [R31]

Reflux

Symptom and specific features	Possible cancer	Recommendation
Reflux (55 and over)	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55

		and over with reflux and nausea or vomiting. [new 2015] [1.2.10] [R40]
Reflux with weight loss (55 and over)	Oesophageal	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people: <ul style="list-style-type: none"> aged 45 and over with dysphagia or aged 55 and over with weight loss and upper abdominal pain, or reflux or dyspepsia. [new 2015] [1.2.1] [R29]
Reflux with weight loss (55 and over)	Stomach	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or are aged 55 and over with upper abdominal pain, reflux or dyspepsia. [new 2015] [1.2.8] [R38]

Bleeding

See also [Primary care investigations](#) for faecal occult blood

Bleeding, bruising or petechiae

Symptom and specific features	Possible cancer	Recommendation
Bleeding - unexplained (children and young people)	Leukaemia	Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms: <ul style="list-style-type: none"> pallor persistent fatigue unexplained fever unexplained persistent infection generalised lymphadenopathy persistent or unexplained bone pain unexplained bruising unexplained bleeding. [new 2015] [1.8.2] [R106]
Petechiae (children and young people)	Leukaemia	Refer children and young people for immediate specialist assessment for leukaemia if they have unexplained

		petechiae or hepatosplenomegaly. [new 2015] [1.8.3] [R107]
Bruising - unexplained (children and young people)	Leukaemia	<p>Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms:</p> <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent infection • generalised lymphadenopathy • persistent or unexplained bone pain • unexplained bruising • unexplained bleeding. [new 2015] [1.8.2] [R106]
Bruising or petechiae - unexplained (adults)	Leukaemia	<p>Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms:</p> <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]

Haematemesis

Symptom and specific features	Possible cancer	Recommendation
Haematemesis	Oesophageal	Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people with haematemesis. [new 2015] [1.2.2] [R30]

Haematuria

Symptom and specific features	Possible cancer	Recommendation
Haematuria – visible)	Wilm's Tumour	Consider very urgent referral (for an

(children)		appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. [new 2015] [1.10.4] [R120]
Haematuria – visible (45 and over)	Bladder	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have unexplained visible haematuria without urinary tract infection. [new 2015] [1.5.4] [R75]
Haematuria – visible (45 and over)	Renal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have unexplained visible haematuria. [new 2015] [1.5.8] [R79]
Haematuria – visible with vaginal discharge (women 55 and over)	Endometrial	Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with unexplained symptoms of vaginal discharge who: <ul style="list-style-type: none"> • are presenting with these symptoms for the first time or • have a high platelet count or • report haematuria. [new 2015] [1.4.12] [R67]
Haematuria – visible with low haemoglobin or high platelet count or high blood glucose (women 55 and over)	Endometrial	Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 years and over with visible haematuria and any of the following: <ul style="list-style-type: none"> • low haemoglobin levels or • high platelet count or • high blood glucose levels. [new 2015] [1.4.13] [R68]
Haematuria – visible (men)	Prostate	Consider a prostate-specific antigen (PSA) test and digital rectal examination to assess for prostate cancer in men with: <ul style="list-style-type: none"> • any lower urinary tract symptoms, such as nocturia, urinary frequency, hesitancy, urgency or retention or • erectile dysfunction or • visible haematuria. [new 2015] [1.5.2] [R73] See also primary care investigations
Haematuria – non-	Bladder	Refer people using a suspected

visible with dysuria or a raised white cell count (60 and over)		cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 60 and over and have unexplained non-visible haematuria and either dysuria or a raised white cell count on a blood test. [new 2015] [1.5.6] [R77]
Haematuria – visible (45 and over)	Renal or bladder	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. [new 2015] [1.5.5] [R76]</p> <p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. [new 2015] [1.5.9] [R80]</p>

Haemoptysis

Symptom and specific features	Possible cancer	Recommendation
Haemoptysis (40 and over)	Lung	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:</p> <ul style="list-style-type: none"> • have chest X-ray findings that suggest lung cancer or • are aged over 55 with haemoptysis or • are aged 40–55, smoke or have smoked in the past and have haemoptysis or • are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> – cough – fatigue – shortness of breath – chest pain – weight loss – appetite loss. [new 2015]

[1.1.1] [R19]

Post-menopausal bleeding

Symptom and specific features	Possible cancer	Recommendation
Post-menopausal bleeding (women 55 and over)	Endometrial	Refer women using a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer if they are aged 55 and over with post-menopausal bleeding (unexplained vaginal bleeding more than 12 months after menstruation has stopped because of the menopause). [new 2015] [1.4.10] [R65]
Post-menopausal bleeding (women under 55)	Endometrial	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer in women aged under 55 with post-menopausal bleeding. [new 2015] [1.4.11] [R66]

Rectal bleeding

Symptom and specific features	Possible cancer	Recommendation
Rectal bleeding (50 and over)	Colorectal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 50 and have unexplained rectal bleeding. [new 2015] [1.3.1] [R44]
Rectal bleeding with abdominal pain or change in bowel habit or weight loss or iron-deficiency anaemia (Under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]
Rectal bleeding	Colorectal	Offer a digital rectal examination to people In patients with unexplained symptoms related to the lower gastrointestinal tract. [new 2015]

[1.3.9] [Ra3]

Vulval bleeding

Symptom and specific features	Possible cancer	Recommendation
Vulval bleeding (women)	Vulval	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for vulval cancer in women with an unexplained vulval lump, ulceration or bleeding. [new 2015] [1.4.15] [R70]

Gynaecological symptoms

See also:

- [Bleeding](#) for vulval bleeding and post-menopausal (vaginal) bleeding
- [Lumps and masses](#) for vaginal lump
- [Skin or surface symptoms](#) for vulval lump and vulval ulceration

Gynaecological examination findings

Symptom and specific features	Possible cancer	Recommendation
Gynaecological examination	Cervical	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for women if the appearance of their cervix is consistent with cervical cancer. [new 2015] [1.4.14] [R69]

Vaginal discharge

Symptom and specific features	Possible cancer	Recommendation
Vaginal discharge with first presentation or high platelet count or haematuria (women 55 and over)	Endometrial	Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with unexplained symptoms of vaginal discharge who: <ul style="list-style-type: none"> are presenting with these symptoms for the first time or have a high platelet count or report haematuria. [new 2015] [1.4.12] [R67]

Lumps or masses

See also:

- [Abdominal symptoms](#) for abdominal and pelvic lumps or masses
- [Skin or surface symptoms](#) for penile mass and vulval lump.

Lumps and masses

Symptom and specific features	Possible cancer	Recommendation
Breast lump (30 and over)	Breast	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 30 and over and have a breast lump with or without pain. [new 2015] [1.3.11] [R53]
Axillary lump (30 and over)	Breast	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer in people aged 30 years and over with an unexplained lump in the axilla. [new 2015] [1.3.13] [R55]

Anal mass	Anal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for anal cancer in people with an unexplained anal mass or unexplained anal ulceration. [new 2015] [1.3.10] [R52]
Neck lump (45 and over)	Laryngeal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with an unexplained lump in the neck. [new 2015] [1.7.2] [R97]
Lip or oral cavity lump	Oral	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a lump on the lip or in the oral cavity that has been assessed by a dental surgeon to be consistent with oral cancer. [new 2015] [1.7.4] [R99]
Neck lump	Oral	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with a persistent and unexplained lump in the neck. [new 2015] [1.7.5] [R101]
Lip or oral cavity lump	Oral	Consider an urgent referral (for an appointment within 2 weeks) for assessment for oral cancer by the community dental service in people with an unexplained lump on the lip or in the oral cavity that has not been assessed by a dental surgeon. [new 2015] [1.7.3] [R98]
Thyroid lump	Thyroid	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for thyroid cancer in people with an unexplained thyroid lump. [new 2015] [1.7.7] [R102]
Vaginal lump (women)	Vaginal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for vaginal cancer in women with an unexplained palpable mass in or at the entrance to the vagina. [new 2015] [1.4.16] [R71]
Bone swelling (children and young adults)	Bone sarcoma	Consider an urgent direct access X-ray (within 2 weeks) to assess for bone sarcoma in children and young adults with unexplained bone swelling or pain. [new 2015] [1.9.1] [R113]
Soft tissue lump increasing in size	Soft Tissue Sarcoma	Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for soft tissue sarcoma in people with an unexplained lump that

		is increasing in size. [new 2015] [1.9.3] [R115]
--	--	--

Lymphadenopathy

Symptom and specific features	Possible cancer	Recommendation
Lymphadenopathy (children and young people)	Leukaemia	<p>Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms:</p> <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent infection • generalised lymphadenopathy • persistent or unexplained bone pain • unexplained bruising • unexplained bleeding. [new 2015] [1.8.2] [R106]
Lymphadenopathy (adults)	Leukaemia	<p>Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms:</p> <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]
Lymphadenopathy	Hodgkin's Lymphoma	<p>Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]</p>
Lymphadenopathy	Non-Hodgkin's Lymphoma	<p>Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in</p>

		people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]
Lymphadenopathy (40 and over)	Lung	Consider an urgent full blood count and chest X-ray (within 2 weeks) to assess for lung cancer in people aged 40 and over with any of the following: <ul style="list-style-type: none"> • finger clubbing or • supraclavicular lymphadenopathy or persistent cervical lymphadenopathy or • chest signs compatible with lung cancer. [new 2015] [1.1.5] [R23]

Oral lesions

Symptom and specific features	Possible cancer	Recommendation
Ulceration in the oral cavity	Oral	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with unexplained ulceration in the oral cavity lasting for more than 14 days. [new 2015] [1.7.5] [R100]

Neurological symptoms

Symptom and specific features	Possible cancer	Recommendation
Newly abnormal cerebellar or other central neurological function (children and young adults)	Brain and central nervous system	Consider very urgent referral (for an appointment within 48 hours) for suspected brain or central nervous system cancer in children and young adults with newly abnormal cerebellar or other central neurological function. [new 2015] [1.7.9] [R104]
Loss of central neurological function (progressive or sub-acute) (adults)	Brain and central nervous system	Consider an urgent direct access MRI scan of the brain (within 2 weeks) to assess for brain or central nervous system cancer in adults with progressive, sub-acute loss of central neurological function. [new 2015] [1.7.8] [R103]

Pain

See also:

- [Abdominal symptoms](#) for abdominal or pelvic pain
- [Respiratory symptoms](#) for chest pain
- [Skeletal symptoms](#) for bone and back pain.

Symptom and specific features	Possible cancer	Recommendation
Alcohol-induced lymph node pain	Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]

Respiratory symptoms**Chest infection**

Symptom and specific features	Possible cancer	Recommendation
Chest infection - persistent or recurrent	Lung	Offer an urgent chest X-ray (within 2 weeks) to assess for lung cancer in people with either: <ul style="list-style-type: none"> thrombocytosis or persistent or recurrent chest infection. [new 2015] [1.1.6] [R24]

Chest pain

Symptom and specific features	Possible cancer	Recommendation
Chest pain with haemoptysis (non-smoker 40–55)	Lung	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they: <ul style="list-style-type: none"> have chest X-ray findings that suggest lung cancer or are aged over 55 with haemoptysis or are aged 40–55, smoke or have smoked in the past and have haemoptysis or are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.1] [R19]
Chest pain (smoker or past smoker 40 and over)	Lung or mesothelioma	Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms: <ul style="list-style-type: none"> cough fatigue shortness of breath

		<ul style="list-style-type: none"> • chest pain • weight loss • appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.7] [R25]
Chest pain with cough or fatigue or shortness of breath or weight loss or appetite loss (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2014] [1.1.3] [R21] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26]
Chest pain (non-smoker 40 and over)	Lung or mesothelioma	Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:

		<ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4] [R22] <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27]
--	--	---

Cough

Symptom and specific features	Possible cancer	Recommendation
Cough with haemoptysis (non-smoker 40–55)	Lung	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:</p> <ul style="list-style-type: none"> • have chest X-ray findings that suggest lung cancer or • are aged over 55 with haemoptysis or • are aged 40–55, smoke or have smoked in the past and have haemoptysis or • are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> – cough – fatigue – shortness of breath – chest pain – weight loss – appetite loss. [new 2015] [1.1.1] [R19]

Chest pain (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 years and over who smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.7] [R25]
Cough with fatigue or shortness of breath or chest pain or weight loss or appetite loss (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2014] [1.1.3] [R21] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath

		<ul style="list-style-type: none"> • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26]
Cough (non-smoker 40 and over)	Lung or Mesothelioma	<p>Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4] [R22] <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27]

Respiratory examination findings

Symptom and specific features	Possible cancer	Recommendation
Chest signs or finger clubbing	Mesothelioma	<p>Offer an urgent chest X-ray (within 2 weeks) to assess for mesothelioma in people aged 40 and over with either:</p> <ul style="list-style-type: none"> • finger clubbing or • chest signs compatible with pleural disease. [new 2015] [1.1.10] [R28]
Chest signs or finger clubbing	Lung	<p>Consider an urgent full blood count and chest X-ray (within 2 weeks) to assess for lung cancer in people aged 40 and over with any of the following:</p> <ul style="list-style-type: none"> • finger clubbing or • supraclavicular lymphadenopathy or persistent cervical lymphadenopathy or

		<ul style="list-style-type: none"> chest signs compatible with lung cancer. [new 2015] [1.1.5] [R23]
--	--	--

Hoarseness

Symptom and specific features	Possible cancer	Recommendation
Hoarseness – persistent and unexplained (45 and over)	Laryngeal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with persistent unexplained hoarseness. [new 2015] [1.7.1] [R96]

Shortness of breath

Symptom and specific features	Possible cancer	Recommendation
Shortness of breath with haemoptysis (40–50 years)	Lung	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:</p> <ul style="list-style-type: none"> have chest X-ray findings that suggest lung cancer or are aged over 55 with haemoptysis or are aged 40–55, smoke or have smoked in the past and have haemoptysis or are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.1] [R19]
Shortness of breath (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> cough fatigue shortness of breath

		<ul style="list-style-type: none"> • chest pain • weight loss • appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 years and over who smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.7] [R25]
Shortness of breath with cough, fatigue, chest pain, weight loss or appetite loss. (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2014] [1.1.3] [R21] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26]
Shortness of breath (non-smoker 40 and over)	Lung or mesothelioma	Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained

		<p>symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4] [R22] <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27]
Shortness of breath with lymphadenopathy	Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]
Shortness of breath with lymphadenopathy or splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]

Skeletal symptoms

Back pain

Symptom and specific features	Possible cancer	Recommendation
--------------------------------------	------------------------	-----------------------

Back pain (60 and older)	Myeloma	Offer a full blood count, calcium level and test for plasma viscosity or erythrocyte sedimentation rate to assess for myeloma in people 60 and older with persistent bone pain, particularly back pain or unexplained fracture. [new 2015] [1.8.4] [R108] See also primary care investigations
Bone pain with weight loss (60 and older)	Pancreatic	Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms: <ul style="list-style-type: none"> • diarrhoea • back pain • abdominal pain • nausea or vomiting • constipation • new-onset diabetes. [new 2015] [1.2.5] [R33]

Bone pain

Symptom and specific features	Possible cancer	Recommendation
Bone pain (children and young people)	Leukaemia	Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms: <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent infection • generalised lymphadenopathy • persistent or unexplained bone pain • unexplained bruising • unexplained bleeding. [new 2015] [1.8.2] [R106]
Bone pain (60 and older)	Myeloma	Offer a full blood count, calcium level and test for plasma viscosity or erythrocyte sedimentation rate to assess for myeloma in people 60 and older with persistent bone pain, particularly back pain or unexplained fracture. [new 2015] [1.8.4] [R108]

		See also primary care investigations
Bone pain (children and young people)	Bone sarcoma	Consider an urgent direct access X-ray (within 2 weeks) to assess for bone sarcoma in children and young adults with unexplained bone swelling or pain. [new 2015] [1.9.1] [R113]

Fracture

Symptom and specific features	Possible cancer	Recommendation
Fracture - unexplained (60 and older)	Myeloma	Offer a full blood count, calcium level and test for plasma viscosity or erythrocyte sedimentation rate to assess for myeloma in people 60 years and older with persistent bone pain, particularly back pain or unexplained fracture. [new 2015] [1.8.4] [R108] See also primary care investigations

Skin or surface symptoms

See also [Lumps and masses](#) for oral lesions.

Symptom and specific features	Possible cancer	Recommendation
Nipple changes of concern, including discharge and retraction (50 and over)	Breast	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 50 and over with any of the following symptoms in one nipple only: <ul style="list-style-type: none"> • discharge • retraction • other changes of concern. [new 2015] [1.3.12] [R54]

Penile lesion - persistent (men)	Penile	Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a persistent penile lesion after treatment for a sexually transmitted infection has been completed. [new 2015] [1.5.13] [R88]
Penile ulcerated lesion (men)	Penile	Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a penile mass or ulcerated lesion, and sexually transmitted infection has been excluded as a cause. [new 2015] [1.5.12] [R87]
Penile symptoms affecting the foreskin or glans - persistent (men)	Penile	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer in men with unexplained or persistent symptoms affecting the foreskin or glans. [new 2015] [1.5.12] [R89]
Dermatoscopy findings	Melanoma	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if dermatoscopy suggests malignant melanoma of the skin. [new 2015] [1.6.1] [R90]
Pigmented skin lesion	Melanoma	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for malignant melanoma if they present with a suspicious pigmented skin lesion that has a weighted 7-point checklist score of 3 or more. [new 2015] [1.6.2] [Ra2]
Anal ulceration	Anal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for anal cancer in people with an unexplained anal mass or unexplained anal ulceration. [new 2015] [1.3.10] [R52]
Skin lesion	Basal cell carcinoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for basal cell carcinoma if there is concern that a delay may have an unfavourable impact, because of factors such as lesion site or size. [new 2015] [1.6.4] [R94]

Skin lesion	Basal cell carcinoma	Consider routine referral for people with a skin lesion that raises the suspicion of a basal cell carcinoma ⁵ . [new 2015] [1.6.4] [R93] GPs should only excise suspected basal cell carcinomas in accordance with the NICE guideline on Improving outcomes for people with skin tumours including melanoma . [new 2015] [1.6.6] [R95]
Skin lesion	Squamous cell carcinoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a skin lesion that raises the suspicion of squamous cell carcinoma. [new 2015] [1.6.3] [R92]
Vulval ulceration	Vulval	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for vulval cancer in women with an unexplained vulval lump, ulceration or bleeding. [new 2015] [1.4.15] [R70]

Urological symptoms

Dysuria

Symptom and specific features	Possible cancer	Recommendation
Dysuria with weight loss (60 and over)	Bladder	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 60 and over and have unexplained non-visible haematuria and either dysuria or a raised white cell count on a blood test. [new 2015] [1.5.6] [R77]

Erectile dysfunction

Symptom and specific features	Possible cancer	Recommendation
Erectile dysfunction	Prostate	Consider a prostate-specific antigen (PSA) test and digital rectal examination to assess for prostate cancer in men with: <ul style="list-style-type: none"> any lower urinary tract symptoms,

⁵ Typical features of basal cell carcinoma include: an ulcer with a raised rolled edge; prominent fine blood vessels around a lesion; or a nodule on the skin (particularly pearly or waxy nodules).

		<p>such as nocturia, urinary frequency, hesitancy, urgency or retention or</p> <ul style="list-style-type: none"> • erectile dysfunction or • visible haematuria. [new 2015] [1.5.2] [R73] <p>See also primary care investigations</p>
--	--	---

Testicular symptoms

Symptom and specific features	Possible cancer	Recommendation
Enlargement or change in shape or texture of the testis - non-painful	Testicular	Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for testicular cancer if they have a non-painful enlargement or change in shape or texture of the testis. [new 2015] [1.5.10] [R85]
Testicular symptoms - unexplained or persistent	Testicular	Consider a direct access ultrasound scan as part of clinical reassessment for testicular cancer in men with unexplained or persistent testicular symptoms. [new 2015] [1.5.11] [R86]

Urinary tract symptoms

Symptom and specific features	Possible cancer	Recommendation
Urinary tract infection - unexplained and recurrent or persistent	Bladder	Consider referral for bladder cancer in people aged 60 years and over with recurrent or persistent urinary tract infection that is unexplained. [new 2015] [1.5.7] [R78]
Urinary urgency and/or frequency	Ovarian	<p>Carry out tests in primary care if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month:</p> <ul style="list-style-type: none"> • persistent abdominal distension (women often refer to this as 'bloating') • feeling full (early satiety) and/or loss of appetite • pelvic or abdominal pain • increased urinary urgency and/or frequency. [1.4.2] [R57] <p>Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61]</p>

		See also primary care investigations
Lower urinary tract symptoms, such as nocturia, urinary frequency, hesitancy, urgency or retention	Prostate	<p>Consider a prostate-specific antigen (PSA) test and digital rectal examination to assess for prostate cancer in men with:</p> <ul style="list-style-type: none"> any lower urinary tract symptoms, such as nocturia, urinary frequency, hesitancy, urgency or retention or erectile dysfunction or visible haematuria. [new 2015] [1.5.2] [R73] <p>See also primary care investigations</p>

Non-specific features of cancer

Appetite loss or early satiety

Symptom and specific features	Possible cancer	Recommendation
Appetite loss	Multiple, including lung, oesophageal, stomach, colorectal, pancreatic, bladder or renal	<p>For people with unexplained appetite loss, which is a symptom of several cancers including lung, oesophageal, stomach, colorectal, pancreatic, bladder and renal cancer:</p> <ul style="list-style-type: none"> carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely and offer urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks). [new 2015] [1.11.4] [R124]
Appetite loss with haemoptysis (non-smoker 40-50)	Lung	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:</p> <ul style="list-style-type: none"> have chest X-ray findings that suggest lung cancer or are aged over 55 with haemoptysis or are aged 40–55, smoke or have smoked in the past and have haemoptysis or are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following

		<p>symptoms:</p> <ul style="list-style-type: none"> – cough – fatigue – shortness of breath – chest pain – weight loss – appetite loss. [new 2015] [1.1.1] [R19]
Appetite loss (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 years and over who smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.7] [R25]
Appetite loss with cough or fatigue or shortness of breath or chest pain or weight loss (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2014] [1.1.3] [R21]

		<p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26]
--	--	--

Appetite loss (non-smoker 40 and over)	Lung or Mesothelioma	<p>Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4] [R22] <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27]
Appetite loss (persistent or frequent) (women especially 50 and over)	Ovarian	<p>Carry out tests in primary care if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month:</p> <ul style="list-style-type: none"> • persistent abdominal distension (women often refer to this as 'bloating') • feeling full (early satiety) and/or loss of appetite • pelvic or abdominal pain • increased urinary urgency and/or frequency. [1.4.2] [R57] <p>Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61]</p> <p>See also primary care investigations</p>
Appetite loss with weight loss	Stomach	<p>Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who:</p> <ul style="list-style-type: none"> • also have appetite loss or • are aged under 55 with dyspepsia

		or upper abdominal pain lasting 2 weeks or more or <ul style="list-style-type: none"> are aged 55 and over with nausea or vomiting. [new 2015] [1.2.9] [R39]
--	--	---

Deep vein thrombosis

Symptom and specific features	Possible cancer	Recommendation
Deep vein thrombosis	Multiple, including uro-genital, breast, colorectal or lung	For people with deep vein thrombosis, which is associated with several cancers including uro-genital, breast, colorectal and lung cancer: <ul style="list-style-type: none"> carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely and consider urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks). [new 2015] [1.11.5] [R125]

Fatigue

Symptom and specific features	Possible cancer	Recommendation
Fatigue – persistent (children and young people)	Leukaemia	Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms: <ul style="list-style-type: none"> pallor persistent fatigue unexplained fever unexplained persistent infection generalised lymphadenopathy persistent or unexplained bone pain unexplained bruising unexplained bleeding. [new 2015] [1.8.2] [R106]
Fatigue – persistent (adults)	Leukaemia	Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms: <ul style="list-style-type: none"> pallor persistent fatigue

		<ul style="list-style-type: none"> • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]
Fatigue with haemoptysis (non-smoker 40–55)	Lung	<p>Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:</p> <ul style="list-style-type: none"> • have chest X-ray findings that suggest lung cancer or • are aged over 55 with haemoptysis or • are aged 40–55, smoke or have smoked in the past and have haemoptysis or • are aged 40–55, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> – cough – fatigue – shortness of breath – chest pain – weight loss – appetite loss. [new 2015] [1.1.1] [R19]
Fatigue (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 years and over who</p>

		<p>smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.7] [R25]
Fatigue with cough or shortness of breath or chest pain or weight loss or appetite loss (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2014] [1.1.3] [R21] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26]
Fatigue (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4]

		<p>[R22]</p> <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27]
Fatigue (women)	Ovarian	<p>Consider carrying out tests in primary care if a woman reports unexplained weight loss, fatigue or changes in bowel habit. [1.4.3] [R58]</p> <p>Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61]</p> <p>See also primary care investigations</p>

Fever

See also [Respiratory symptoms](#) for chest infection.

Symptom and specific features	Possible cancer	Recommendation
Fever – unexplained (children and young people)	Leukaemia	<p>Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms:</p> <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent infection • generalised lymphadenopathy • persistent or unexplained bone pain • unexplained bruising • unexplained bleeding. [new 2015] [1.8.2] [R106]

Fever – unexplained (adults)	Leukemia	Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms: <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]
Fever with unexplained lymphadenopathy	Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]
Fever with unexplained lymphadenopathy or splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]

Diabetes

Symptom and specific features	Possible cancer	Recommendation
Diabetes - new onset (60 and over)	Pancreatic	Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms: <ul style="list-style-type: none"> • diarrhoea • back pain

		<ul style="list-style-type: none">• abdominal pain• nausea or vomiting• constipation• new-onset diabetes. [new 2015] [1.2.5] [R33]
--	--	--

Night sweats

Symptom and specific features	Possible cancer	Recommendation
Night sweats with unexplained lymphadenopathy	Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]
Night sweats with lymphadenopathy or splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]

Pallor

Symptom and specific features	Possible cancer	Recommendation
Pallor (children and young people)	Leukaemia	Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young people with any of the following symptoms: <ul style="list-style-type: none"> • pallor • persistent fatigue • unexplained fever • unexplained persistent infection • generalised lymphadenopathy • persistent or unexplained bone pain • unexplained bruising • unexplained bleeding. [new 2015] [1.8.2] [R106]
Pallor (adults)	Leukaemia	Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms: <ul style="list-style-type: none"> • pallor

		<ul style="list-style-type: none"> • persistent fatigue • unexplained fever • unexplained persistent or recurrent infection • generalised lymphadenopathy • unexplained bruising • unexplained bleeding • unexplained petechiae • hepatosplenomegaly. [new 2015] [1.8.1] [R105]
--	--	--

Parental concern

Symptom and specific features	Possible cancer	Recommendation
Parental concern	Childhood cancer	Take into account the insight and knowledge of parents and carers when considering urgent referral for a child or young person. [2015] [1.11.1] [R121]

Parental concern or anxiety	Childhood cancer	Consider referral for children if their parent or carer has persistent concern or anxiety about the child's symptoms, even if the symptoms are most likely to have a benign cause. [2015] [1.11.2] [R122]
-----------------------------	------------------	--

Pruritus

Symptom and specific features	Possible cancer	Recommendation
Pruritus with lymphadenopathy or splenomegaly	Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]
Pruritus with lymphadenopathy or splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]

Weight loss

Symptom and specific features	Possible cancer	Recommendation
Weight loss – unexplained	Colorectal, gastro-oesophageal, lung, prostate, pancreatic or urological cancer	For people with unexplained weight loss, which is a symptom of several cancers including colorectal, gastro-oesophageal, lung, prostate, pancreatic and urological cancer: <ul style="list-style-type: none"> carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely and offer urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks) [new 2015] [1.11.3] [R123]

Weight loss – unexplained with abdominal pain (over 40)	Colorectal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 40 with unexplained weight loss and abdominal pain. [new 2015] [1.3.5] [R48]
Weight loss with rectal bleeding (under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]
Weight loss without rectal bleeding	Colorectal	Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who: <ul style="list-style-type: none"> • have abdominal pain or • have weight loss or • are aged under 60 years and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.6] [R49] See also primary care investigations
Weight loss – unexplained	Colorectal	Offer a digital rectal examination to people In patients with unexplained symptoms related to the lower gastrointestinal tract. [new 2015] [1.3.9] [Ra3]
Weight loss with haemoptysis (non-smoker 40–55)	Lung	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they: <ul style="list-style-type: none"> • have chest X-ray findings that suggest lung cancer or • are aged over 55 with haemoptysis or • are aged 40–55, smoke or have smoked in the past and have haemoptysis or

		<ul style="list-style-type: none"> are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.1] [R19]
Weight loss (smoker or past smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.2] [R20] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 years and over who smoke or have smoked in the past with any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.7] [R25] <p>See also primary care investigations</p>
Weight loss with cough, fatigue, shortness of breath, chest pain or appetite loss (non-smoker 40 and over)	Lung or mesothelioma	<p>Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:</p> <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain

		<ul style="list-style-type: none"> • weight loss • appetite loss. [new 2014] [1.1.3] [R21] <p>Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.8] [R26] <p>See also primary care investigations</p>
Weight loss (non-smoker 40 and over)	Lung or Mesothelioma	<p>Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.4] [R22] <p>Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:</p> <ul style="list-style-type: none"> • cough • fatigue • shortness of breath • chest pain • weight loss • appetite loss. [new 2015] [1.1.9] [R27] <p>See also primary care investigations</p>
Weight loss with lymphadenopathy or splenomegaly	Hodgkin's Lymphoma	<p>Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly</p>

		fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. [new 2015] [1.8.9] [R112]
Weight loss with lymphadenopathy or splenomegaly	Non-Hodgkin's Lymphoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. [new 2015] [1.8.8] [R111]
Weight loss with upper abdominal pain or reflux or dyspepsia (55 and over)	Oesophageal	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people: <ul style="list-style-type: none"> aged 45 and over with dysphagia or aged 55 and over with weight loss and upper abdominal pain, or reflux or dyspepsia. [new 2015] [1.2.1] [R29]
Weight loss with nausea or vomiting (55 and over)	Oesophageal	Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people aged 55 or over with: <ul style="list-style-type: none"> weight loss and nausea or vomiting or reflux and nausea or vomiting or upper abdominal pain and raised platelet count. [new 2015] [1.2.3] [R31]
Weight loss with upper abdominal pain and nausea or vomiting (40 and over) or Weight loss with upper abdominal pain or reflux or dyspepsia (55 and over)	Stomach	Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who: <ul style="list-style-type: none"> are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or are aged 55 and over with upper abdominal pain, reflux or dyspepsia. [new 2015] [1.2.8] [R38]
Weight loss with appetite loss or Weight loss with dyspepsia or upper abdominal pain	Stomach	Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who:

(under 55) or Weight loss with nausea or vomiting (55 and over)		<ul style="list-style-type: none"> • also have appetite loss or • are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more or • are aged 55 and over with nausea or vomiting. [new 2015] [1.2.9] [R39]
Weight loss (women)	Ovarian	<p>Consider carrying out tests in primary care if a woman reports unexplained weight loss, fatigue or changes in bowel habit. [1.4.3] [R58]</p> <p>Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer). [1.4.6] [R61]</p> <p>See also primary care investigations</p>
Weight loss with diarrhoea or back pain or abdominal pain or nausea or vomiting or constipation or new- onset diabetes	Pancreatic	<p>Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms:</p> <ul style="list-style-type: none"> • diarrhoea • back pain • abdominal pain • nausea or vomiting • constipation • new-onset diabetes. [new 2015] [1.2.5] [R33]

Primary care investigations

See also:

- [Abdominal symptoms](#) for abdominal examination findings.
- [Gynaecological symptoms](#) for gynaecological examination findings

Anaemia

Investigation findings and specific features	Possible cancer	Recommendation
Anaemia (iron deficiency) (over 60)	Colorectal	Refer people using a suspected cancer pathway referral (with an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.2] [R45]
Anaemia (iron deficiency) without rectal bleeding (under 60)	Colorectal	Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who: <ul style="list-style-type: none"> • have abdominal pain or • have weight loss or • are aged under 60 years and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.6] [R49] See also primary care investigations
Anaemia (iron deficiency) with rectal bleeding (under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]

Blood test findings

Investigation findings and specific features	Possible cancer	Recommendation
Raised white cell count with non-visible haematuria	Bladder	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 60 and

		over and have unexplained non-visible haematuria and either dysuria or a raised white cell count on a blood test. [new 2015] [1.5.6] [R77]
Anaemia (iron-deficiency) (over 60)	Colorectal	Refer people using a suspected cancer pathway referral (with an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.2] [R45]
Anaemia (iron deficiency) with rectal bleeding (under 50)	Colorectal	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings: <ul style="list-style-type: none"> • abdominal pain • change in bowel habit • weight loss • iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.8] [R51]
Anaemia (iron deficiency) without rectal bleeding (under 60)	Colorectal	Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who: <ul style="list-style-type: none"> • have abdominal pain or • have weight loss or • are aged under 60 years and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). [new 2015] [1.3.6] [R49]
Protein electrophoresis results suggest myeloma	Myeloma	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if the results of protein electrophoresis suggest myeloma. [new 2015] [1.8.7] [R110]
Plasma viscosity or erythrocyte sedimentation rate results and presentation consistent with	Myeloma	Consider very urgent protein electrophoresis (within 48 hours) to assess for myeloma if the plasma viscosity or erythrocyte sedimentation rate results and presentation are consistent with possible myeloma.

possible myeloma		[new 2015] [1.8.6] [R109]
Hypercalcaemia or leucopenia and presentation consistent with possible myeloma	Myeloma	Offer very urgent protein electrophoresis (within 48 hours) to assess for myeloma in people 60 years and older with hypercalcaemia or leucopenia and a presentation that is consistent with possible myeloma. [new 2015] [1.8.5] [Ra1]
Prostate-specific antigen levels increased	Prostate	Refer men using a suspected cancer pathway referral (with an appointment within 2 weeks) for prostate cancer if their PSA levels are above the age-specific reference range. [new 2015] [1.5.3] [R74]
Thrombocytosis	Lung	Offer an urgent chest X-ray (within 2 weeks) to assess for lung cancer in people with either: <ul style="list-style-type: none"> • thrombocytosis or • persistent or recurrent chest infection. [new 2015] [1.1.6] [R24]
Serum CA125 35 IU/ml or greater	Ovarian	If serum CA125 is 35 IU/ml or greater, arrange an ultrasound scan of the abdomen and pelvis. [1.4.7] [R62] For any woman who has normal serum CA125 (less than 35 IU/ml), or CA125 of 35 IU/ml or greater but a normal ultrasound: <ul style="list-style-type: none"> • assess her carefully for other clinical causes of her symptoms and investigate if appropriate • if no other clinical cause is apparent, advise her to return to her GP if her symptoms become more frequent and/or persistent. [1.4.9] [R64]
High platelet count with unexplained vaginal discharge (women 55 and over)	Endometrial	Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with unexplained symptoms of vaginal discharge who: <ul style="list-style-type: none"> • are presenting with these symptoms for the first time or • have a high platelet count or • report haematuria. [new 2015] [1.4.12] [R67]
Low haemoglobin levels or high platelet count or	Endometrial	Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 years and over with

high blood glucose levels, with visible haematuria (women 55 and over)		visible haematuria and any of the following: <ul style="list-style-type: none"> • low haemoglobin levels or • high platelet count or • high blood glucose levels. [new 2015] [1.4.13] [R68]
New-onset diabetes	Pancreatic	Consider an urgent direct access CT scan (within 2 weeks), or urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms: <ul style="list-style-type: none"> • diarrhoea • back pain • abdominal pain • nausea or vomiting • constipation • new-onset diabetes. [new 2015] [1.2.5] [R33]

Faecal tests

Investigation findings and specific features	Possible cancer	Recommendation
Occult blood in faeces	Colorectal	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if tests show occult blood in their faeces. [new 2015] [1.3.7] [R50]

Imaging tests

Investigation findings and specific features	Possible cancer	Recommendation
Chest X-ray findings that suggest lung cancer	Lung	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they: <ul style="list-style-type: none"> • have chest X-ray findings that suggest lung cancer or • are aged over 55 with haemoptysis or • are aged 40–55, smoke or have smoked in the past and have haemoptysis or

		<ul style="list-style-type: none"> are aged 40–55 years, have never smoked and have haemoptysis and at least one of the following symptoms: <ul style="list-style-type: none"> cough fatigue shortness of breath chest pain weight loss appetite loss. [new 2015] [1.1.1] [R19]
Ultrasound findings that suggest ovarian cancer	Ovarian	If the ultrasound suggests ovarian cancer, refer the woman urgently ⁶ for further investigation. [1.4.8] [R63]
X-ray suggests the possibility of bone sarcoma	Bone sarcoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if an X-ray suggests the possibility of bone sarcoma. [new 2015] [1.9.2][R114]
Ultrasound findings that are suggestive of soft tissue sarcoma	Soft tissue sarcoma	Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if they have ultrasound findings that are suggestive of soft tissue sarcoma or if ultrasound findings are uncertain and clinical concern persists. [new 2015] [1.9.4] [R116]

Jaundice

Investigation findings and specific features	Possible cancer	Recommendation
Jaundice (40 and over)	Pancreatic	Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for pancreatic cancer if they are aged 40 and over and have jaundice. [new 2015] [1.2.4] [R32]

Ocular examination

Investigation findings and specific features	Possible cancer	Recommendation

⁶ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

Absent red reflex	Retinoblastoma	Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex. [new 2015] [1.10.2] [R118]
-------------------	----------------	--

Recommendations organised by site of cancer

The following guidance is based on the best available evidence. The [full guideline](#) [\[hyperlink to be added for final publication\]](#) gives details of the methods and the evidence used to develop the guidance.

1.1 *Lung and pleural cancers*

Lung cancer

- 1.1.1 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:
- have chest X-ray findings that suggest lung cancer **or**
 - are aged over 55 with haemoptysis **or**
 - are aged 40-55, smoke or have smoked in the past, and have haemoptysis **or**
 - are aged 40-55, have never smoked and have haemoptysis and at least one of the following symptoms:
 - cough
 - fatigue
 - shortness of breath
 - chest pain
 - weight loss
 - appetite loss. **[new 2015]** [R19]
- 1.1.2 Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:
- cough
 - fatigue
 - shortness of breath
 - chest pain
 - weight loss
 - appetite loss. **[new 2015]** [R20]

1.1.3 Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. **[new 2015]** [R21]

1.1.4 Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. **[new 2015]** [R22]

1.1.5 Consider an urgent full blood count and chest X-ray (within 2 weeks) to assess for lung cancer in people aged 40 and over with any of the following:

- finger clubbing or
- supraclavicular lymphadenopathy or persistent cervical lymphadenopathy or
- chest signs compatible with lung cancer. **[new 2015]** [R23]

1.1.6 Offer an urgent chest X-ray (within 2 weeks) to assess for lung cancer in people with either:

- thrombocytosis or
- persistent or recurrent chest infection. **[new 2015]** [R24]

Mesothelioma

1.1.7 Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. **[new 2015]** [R25]

1.1.8 Offer a full blood count and chest X-ray to assess for mesothelioma in people aged 40 and over who have never smoked and have 2 or more of the following symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. **[new 2015]** [R26]

1.1.9 Offer a full blood count to assess for mesothelioma in people aged 40 and over who have never smoked and have any of the following symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. **[new 2015]** [R27]

1.1.10 Offer an urgent chest X-ray (within 2 weeks) to assess for mesothelioma in people aged 40 and over with either:

- finger clubbing or
- chest signs compatible with pleural disease. **[new 2015]** [R28]

1.2 *Upper gastro-intestinal tract cancers*

Oesophageal cancer

1.2.1 Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people:

- aged 45 and over with dysphagia **or**
- aged 55 and over with weight loss and upper abdominal pain or reflux or dyspepsia. **[new 2015]** [R29]

1.2.2 Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people with haematemesis. **[new 2015]** [R30]

1.2.3 Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people aged 55 or over with:

- weight loss and nausea or vomiting or
- reflux and nausea or vomiting or
- upper abdominal pain and raised platelet count. **[new 2015]** [R31]

Pancreatic cancer

- 1.2.4 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for pancreatic cancer if they are aged 40 and over and have jaundice. **[new 2015]** [R32]
- 1.2.5 Consider an urgent direct access CT scan (within 2 weeks), or an urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms:
- diarrhoea
 - back pain
 - abdominal pain
 - nausea or vomiting
 - constipation
 - new-onset diabetes. **[new 2015]** [R33]

Stomach cancer

- 1.2.6 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with an upper abdominal mass consistent with stomach cancer. **[new 2015]** [R36]
- 1.2.7 Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with dysphagia. **[new 2015]** [R37]
- 1.2.8 Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who:
- are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or
 - are aged 55 and over with upper abdominal pain, reflux or dyspepsia. **[new 2015]** [R38]

1.2.9 Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who:

- also have appetite loss **or**
- are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more **or**
- are aged 55 and over with nausea or vomiting. **[new 2015]** [R39]

1.2.10 Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with reflux and nausea or vomiting. **[new 2015]** [R40]

1.2.11 Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with upper abdominal pain and raised platelet counts. **[new 2015]** [R41]

Gall bladder cancer

1.2.12 Consider urgent direct access ultrasound (within 2 weeks) to assess for gall bladder cancer in people with an upper abdominal mass consistent with an enlarged gall bladder. **[new 2015]** [R42]

Liver cancer

1.2.13 Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for liver cancer in people with an upper abdominal mass consistent with an enlarged liver. **[new 2015]** [R43]

1.3 *Lower gastrointestinal tract cancers*

Colorectal cancer

1.3.1 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 50 and have unexplained rectal bleeding. **[new 2015]** [R44]

1.3.2 Refer people using a suspected cancer pathway referral (with an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained iron-deficiency anaemia

(haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). **[new 2015]** [R45]

- 1.3.3 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained changes in their bowel habit. **[new 2015]** [R46]
- 1.3.4 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people with a rectal or abdominal mass. **[new 2015]** [R47]
- 1.3.5 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 40 with unexplained weight loss and abdominal pain. **[new 2015]** [R48]
- 1.3.6 Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who:
- have abdominal pain **or**
 - have weight loss **or**
 - are aged under 60 and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). **[new 2015]** [R49]
- 1.3.7 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if tests show occult blood in their faeces. **[new 2015]** [R50]
- 1.3.8 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings:
- abdominal pain
 - change in bowel habit

- weight loss
- iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). **[new 2015]** [R51]

1.3.9 Offer a digital rectal examination to people with unexplained symptoms related to the lower gastrointestinal tract. **[new 2015]** [Ra3]

Anal cancer

1.3.10 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for anal cancer in people with an unexplained anal mass or unexplained anal ulceration. **[new 2015]** [R52]

1.3.11 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 30 and over and have a breast lump with or without pain. **[new 2015]** [R53]

1.3.12 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 50 and over with any of the following symptoms in one nipple only:

- discharge
- retraction
- other changes of concern. **[new 2015]** [R54]

1.3.13 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer in people aged 30 and over with an unexplained lump in the axilla. **[new 2015]** [R55]

1.4 Gynaecological cancer

Ovarian cancer

The recommendations in this section are from the NICE guideline on [ovarian cancer](#) (2011).

- 1.4.1 Refer the woman urgently⁷ if physical examination identifies ascites and/or a pelvic or abdominal mass (which is not obviously uterine fibroids).[R56]
- 1.4.2 Carry out tests in primary care (see recommendations 1.4.6 to 1.4.9) if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month:
- persistent abdominal distension (women often refer to this as 'bloating')
 - feeling full (early satiety) and/or loss of appetite
 - pelvic or abdominal pain
 - increased urinary urgency and/or frequency. [R57]

⁷ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

- 1.4.3 Consider carrying out tests in primary care (see recommendations 1.4.6 to 1.4.9) if a woman reports unexplained weight loss, fatigue or changes in bowel habit. [R58]
- 1.4.4 Advise any woman who is not suspected of having ovarian cancer to return to her GP if her symptoms become more frequent and/or persistent. [R59]
- 1.4.5 Carry out appropriate tests for ovarian cancer (see recommendations 1.4.6 to 1.4.9) in any woman of 50 or over who has experienced symptoms within the last 12 months that suggest irritable bowel syndrome (IBS)⁸, because IBS rarely presents for the first time in women of this age. [R60]
- 1.4.6 Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer (see recommendations 1.4.1 to 1.4.5). [R61]
- 1.4.7 If serum CA125 is 35 IU/ml or greater, arrange an ultrasound scan of the abdomen and pelvis. [R62]
- 1.4.8 If the ultrasound suggests ovarian cancer, refer the woman urgently⁹ for further investigation. [R63]
- 1.4.9 For any woman who has normal serum CA125 (less than 35 IU/ml), or CA125 of 35 IU/ml or greater but a normal ultrasound:
- assess her carefully for other clinical causes of her symptoms and investigate if appropriate
 - if no other clinical cause is apparent, advise her to return to her GP if her symptoms become more frequent and/or persistent. [R64]

⁸ See the NICE guideline on [irritable bowel syndrome in adults](#).

⁹ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

Endometrial cancer

- 1.4.10 Refer women using a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer if they are aged 55 and over with post-menopausal bleeding (unexplained vaginal bleeding more than 12 months after menstruation has stopped because of the menopause). **[new 2015]** [R65]
- 1.4.11 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer in women aged under 55 with post-menopausal bleeding. **[new 2015]** [R66]
- 1.4.12 Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with unexplained symptoms of vaginal discharge who:
- are presenting with these symptoms for the first time **or**
 - have a high platelet count **or**
 - report haematuria. **[new 2015]** [R67]
- 1.4.13 Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with visible haematuria and any of the following:
- low haemoglobin levels **or**
 - high platelet count **or**
 - high blood glucose levels. **[new 2015]** [R68]

Cervical cancer

- 1.4.14 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for women if the appearance of their cervix is consistent with cervical cancer. **[new 2015]** [R69]

Vulval cancer

- 1.4.15 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for vulval cancer in women with an unexplained vulval lump, ulceration or bleeding. **[new 2015]** [R70]

Vaginal cancer

- 1.4.16 Consider a suspected cancer pathway referral (with an appointment within 2 weeks) for vaginal cancer in women with an unexplained palpable mass in or at the entrance to the vagina. **[new 2015]** [R71]

1.5 *Urological cancers*

Prostate cancer

- 1.5.1 Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for prostate cancer if their prostate feels malignant on digital rectal examination. **[new 2015]** [R72]
- 1.5.2 Consider a prostate-specific antigen (PSA) test and digital rectal examination to assess for prostate cancer in men with:
- any lower urinary tract symptoms, such as nocturia, urinary frequency, hesitancy, urgency or retention **or**
 - erectile dysfunction **or**
 - visible haematuria. **[new 2015]** [R73]

- 1.5.3 Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for prostate cancer if their PSA levels are above the age-specific reference range. **[new 2015]** [R74]

Bladder cancer

- 1.5.4 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have unexplained visible haematuria without urinary tract infection. **[new 2015]** [R75]
- 1.5.5 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. **[new 2015]** [R76]
- 1.5.6 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 60 and over and have unexplained non-visible haematuria and either dysuria or a raised white cell count on a blood test. **[new 2015]** [R77]
- 1.5.7 Consider referral for bladder cancer in people aged 60 and over with recurrent or persistent urinary tract infection that is unexplained. **[new 2015]** [R78]

Renal cancer

- 1.5.8 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have unexplained visible haematuria. **[new 2015]** [R79]
- 1.5.9 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. **[new 2015]** [R80]

Testicular cancer

- 1.5.10 Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for testicular cancer if they have a non-painful enlargement or change in shape or texture of the testis. **[new 2015]** [R85]
- 1.5.11 Consider a direct access ultrasound scan as part of clinical reassessment for testicular cancer in men with unexplained or persistent testicular symptoms. **[new 2015]** [R86]

Penile cancer

- 1.5.12 Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a penile mass or ulcerated lesion, and sexually transmitted infection has been excluded as a cause. **[new 2015]** [R87]
- 1.5.13 Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a persistent penile lesion after treatment for a sexually transmitted infection has been completed. **[new 2015]** [R88]
- 1.5.14 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer in men with unexplained or persistent symptoms affecting the foreskin or glans. **[new 2015]** [R89]

1.6 *Skin cancers*

Malignant melanoma of the skin

- 1.6.1 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if dermatoscopy suggests malignant melanoma of the skin. **[new 2015]** [R90]
- 1.6.2 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for malignant melanoma if they

present with a suspicious pigmented skin lesion that has a weighted 7-point checklist score of 3 or more. **[new 2015] [Ra2]**

Weighted 7-point checklist

Major features of the lesions (scoring 2 points each):

- change in size
- irregular shape
- irregular colour.

Minor features of the lesions (scoring 1 point each):

- largest diameter 7 mm or more
- inflammation
- oozing
- change in sensation.

Squamous cell carcinoma

- 1.6.3 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a skin lesion that raises the suspicion of squamous cell carcinoma. **[new 2015] [R92]**

Basal cell carcinoma

- 1.6.4 Consider routine referral for people if they have a skin lesion that raises the suspicion of a basal cell carcinoma¹⁰. **[new 2015] [R93]**
- 1.6.5 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for basal cell carcinoma if there is concern that a delay may have an unfavourable impact, because of factors such as lesion site or size. **[new 2015] [R94]**

¹⁰ Typical features of basal cell carcinoma include: an ulcer with a raised rolled edge; prominent fine blood vessels around a lesion; or a nodule on the skin (particularly pearly or waxy nodules).

- 1.6.6 GPs should only excise suspected basal cell carcinomas in accordance with the NICE guideline on [improving outcomes for people with skin tumours including melanoma](#). **[new 2015]** [R95]

1.7 Head and neck cancers

Laryngeal cancer

- 1.7.1 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with persistent unexplained hoarseness. **[new 2015]** [R96]
- 1.7.2 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with an unexplained lump in the neck. **[new 2015]** [R97]

Oral cancer

- 1.7.3 Consider an urgent referral (for an appointment within 2 weeks) for assessment for oral cancer by the community dental service in people with an unexplained lump on the lip or in the oral cavity that has not been assessed by a dental surgeon. **[new 2015]** [R98]
- 1.7.4 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a lump on the lip or in the oral cavity that has been assessed by a dental surgeon to be consistent with oral cancer. **[new 2015]** [R99]
- 1.7.5 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with unexplained ulceration in the oral cavity lasting for more than 14 days. **[new 2015]** [R100]
- 1.7.6 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with a persistent and unexplained lump in the neck. **[new 2015]** [R101]

Thyroid cancer

- 1.7.7 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for thyroid cancer in people with an unexplained thyroid lump. **[new 2015]** [R102]

Brain and central nervous system cancers

- 1.7.8 Consider an urgent direct access MRI scan of the brain (within 2 weeks) to assess for brain or central nervous system cancer in adults with progressive, sub-acute loss of central neurological function. **[new 2015]** [R103]
- 1.7.9 Consider very urgent referral (for an appointment within 48 hours) for suspected brain or central nervous system cancer in children and young adults with newly abnormal cerebellar or other central neurological function. **[new 2015]** [R104]

1.8 *Haematological cancers*

Leukaemia

- 1.8.1 Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms:
- pallor
 - persistent fatigue
 - unexplained fever
 - unexplained persistent or recurrent infection
 - generalised lymphadenopathy
 - unexplained bruising
 - unexplained bleeding
 - unexplained petechiae
 - hepatosplenomegaly. **[new 2015]** [R105]
- 1.8.2 Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young adults with any of the following symptoms:
- pallor

- persistent fatigue
- unexplained fever
- unexplained persistent infection
- generalised lymphadenopathy
- persistent or unexplained bone pain
- unexplained bruising
- unexplained bleeding. **[new 2015]** [R106]

1.8.3 Refer children and young adults for immediate specialist assessment for leukaemia if they have unexplained petechiae or hepatosplenomegaly. **[new 2015]** [R107]

Myeloma

- 1.8.4 Offer a full blood count, blood tests for calcium and plasma viscosity or erythrocyte sedimentation rate to assess for myeloma in people aged 60 and over with persistent bone pain, particularly back pain or unexplained fracture. **[new 2015]** [R108]
- 1.8.5 Offer very urgent protein electrophoresis (within 48 hours) to assess for myeloma in people aged 60 and over with hypercalcaemia or leucopenia and a presentation that is consistent with possible myeloma. **[new 2015]** [Ra1]
- 1.8.6 Consider very urgent protein electrophoresis (within 48 hours) to assess for myeloma if the plasma viscosity or erythrocyte sedimentation rate and presentation are consistent with possible myeloma. **[new 2015]** [R109]
- 1.8.7 Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if the results of protein electrophoresis suggest myeloma. **[new 2015]** [R110]

Non-Hodgkin's lymphoma

- 1.8.8 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When

considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. **[new 2015]** [R111]

Hodgkin's lymphoma

- 1.8.9 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. **[new 2015]** [R112]

1.9 Sarcomas

Bone sarcoma

- 1.9.1 Consider an urgent direct access X-ray (within 2 weeks) to assess for bone sarcoma in children and young adults with unexplained bone swelling or pain. **[new 2015]** [R113]
- 1.9.2 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if an X-ray suggests the possibility of bone sarcoma. **[new 2015]** [R114]

Soft tissue sarcoma

- 1.9.3 Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for soft tissue sarcoma in people with an unexplained lump that is increasing in size. **[new 2015]** [R115]
- 1.9.4 Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if they have ultrasound scan findings that are suggestive of soft tissue sarcoma or if ultrasound findings are uncertain and clinical concern persists. **[new 2015]** [R116]

1.10 Childhood cancers

Neuroblastoma

- 1.10.1 Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]** [R117]

Retinoblastoma

- 1.10.2 Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex. **[new 2015]** [R118]

Wilm's tumour

- 1.10.3 Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]** [R119]
- 1.10.4 Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. **[new 2015]** [R120]

1.11 *Non-site-specific symptoms*

Symptoms of concern in children

- 1.11.1 Take into account the insight and knowledge of parents and carers when considering urgent referral for a child or young person. **[2015]** [R121]
- 1.11.2 Consider referral for children if their parent or carer has persistent concern or anxiety about the child's symptoms, even if the symptoms are most likely to have a benign cause. **[2015]** [R122]

Symptoms of concern in adults

- 1.11.3 For people with unexplained weight loss, which is a symptom of several cancers including colorectal, gastro-oesophageal, lung, prostate, pancreatic and urological cancer:

- carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely **and**
- offer urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks). **[new 2015]** [R123]

1.11.4 For people with unexplained appetite loss, which is a symptom of several cancers including lung, oesophageal, stomach, colorectal, pancreatic, bladder and renal cancer:

- carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely **and**
- offer urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks). **[new 2015]** [R124]

1.11.5 For people with deep vein thrombosis, which is associated with several cancers including uro-genital, breast, colorectal and lung cancer:

- carry out an assessment for additional symptoms, signs or findings that may help to clarify which cancer is most likely **and**
- consider urgent investigation or a suspected cancer pathway referral (for an appointment within 2 weeks). **[new 2015]** [R125]

Recommendations on patient support, safety netting and the diagnostic process

1.12 *Patient information and support*

- 1.12.1 Discuss with people with suspected cancer (and their carers as appropriate, taking account of the need for confidentiality) their preferences for being involved in decision-making about referral options and further investigations (including their potential risks and benefits), and ensure they have the time for this. **[2015]** [R1]
- 1.12.2 When cancer is suspected in a child, discuss the referral decision and information to be given to the child with the parents or carers (and the child if appropriate). **[2015]** [R2]
- 1.12.3 Explain to people who are being referred with suspected cancer that they are being referred to a cancer service, but if appropriate reassure them that most people referred will not have a diagnosis of cancer, and discuss alternative diagnoses with them. **[2015]** [R3]
- 1.12.4 Give the person information on the possible diagnosis (both benign and malignant) in accordance with their wishes for information. Follow current advice on communicating with patients and/or their carers and breaking bad news (see also [Patient experience in adult NHS services](#) (NICE guideline CG138). **[2015]** [R4]
- 1.12.5 The information given to people with suspected cancer and their families and/or carers should cover, among other issues:
- where the person is being referred to
 - how long they will have to wait for the appointment
 - how to obtain further information about the type of cancer suspected or help prior to the specialist appointment
 - who they will be seen by
 - what to expect from the service the person will be attending

- what type of tests will be carried out, and what will happen during diagnostic procedures
- how long it will take to get a diagnosis or test results
- whether they can take someone with them to the appointment
- other sources of support, including those for minority groups.

[2015] [R5]

- 1.12.6 Provide information that is appropriate for the person in terms of language, ability and culture, recognising the potential for different cultural meanings associated with the possibility of cancer, the relative importance of family decision-making and possible unfamiliarity with the concept of support outside the family. **[2015]** [R6]
- 1.12.7 Have information available in a variety of formats on both local and national sources of additional support for people who are being referred with suspected cancer. For more information on information sharing, see section 1.5 in [Patient experience in adult NHS services](#) (NICE guideline CG138). **[2015]** [R7]
- 1.12.8 Reassure people in the safety netting group (see recommendation X) who are concerned that they may have cancer that with their current symptoms their risk of having cancer is low. **[new 2015]** [R8]
- 1.12.9 Explain to people who are being offered safety netting (see recommendation 2.2.1) which symptoms to look out for and when they should return for re-evaluation. It may be appropriate to provide written information about this. **[new 2015]** [R9]
- 1.12.10 When referring a person with suspected cancer to a specialist service, assess their need for continuing support while waiting for their referral appointment. This should include inviting the person to contact their healthcare professional again if they have more concerns or questions before they see a specialist. **[2005]**
- 1.12.11 If the person has additional support needs because of their personal circumstances, inform the specialist (with the person's agreement). **[2005]**

1.13 Safety netting

1.13.1 Consider a review for people with any symptom that is associated with an increased risk of cancer, but who do not meet the criteria for referral. The review may be:

- planned within a time frame agreed with the person or
- patient-initiated if their symptoms recur, persist or worsen, or new symptoms develop or the person continues to be concerned. **[new 2015]** [R10]

1.14 *The diagnostic process*

- 1.14.1 Take part in continuing education, peer review and other activities to improve and maintain clinical consulting, reasoning and diagnostic skills, in order to identify at an early stage people who may have cancer, and to communicate the possibility of cancer to the person. **[2005]** [R11]
- 1.14.2 Discussion with a specialist (for example, by telephone or email) should be considered if there is uncertainty about the interpretation of symptoms and signs, and whether a referral is needed. This may also enable the primary healthcare professional to communicate their concerns and a sense of urgency to secondary healthcare professionals when symptoms are not classical. **[2005]** [R12]
- 1.14.3 Put in place local arrangements to ensure that letters about non-urgent referrals are assessed by the specialist, so that the person can be seen more urgently if necessary. **[2005]** [R13]
- 1.14.4 Put in place local arrangements to ensure that there is a maximum waiting period for non-urgent referrals, in accordance with national targets and local arrangements. **[2005]** [R14]
- 1.14.5 Ensure local arrangements are in place to identify people who miss their appointments so that they can be followed up. **[2005]** [R15]
- 1.14.6 Include all appropriate information in referral correspondence, including whether the referral is urgent or non-urgent. **[2005]** [R16]
- 1.14.7 Use local referral proformas if these are in use. **[2005]** [R17]
- 1.14.8 Once the decision to refer has been made, make sure that the referral is made within 1 working day. **[2005]** [R18]

2 Research recommendations

The Guideline Development Group has made the following recommendations for research, based on its review of evidence, to improve NICE guidance and patient care in the future.

2.1 *Age thresholds in cancer*

Longitudinal studies to identify and quantify factors in adults that are associated with development of specific cancers at a younger age than the norm. They should be designed to inform age thresholds in clinical guidance. The primary outcome should be likelihood ratios and positive predictive values for cancer occurring in younger age groups.

Why this is important

It is recognised that several factors, such as deprivation and co-morbidity, may lead to development of cancer at a younger age. People with these factors could be disadvantaged by the use of age thresholds for referral for suspected cancer.

2.2 *Primary care testing*

Diagnostic accuracy studies of tests accessible to primary care for a given cancer in symptomatic people. Priority areas for research should include tests for people with cough, non-visible haematuria, suspected prostate cancer, suspected pancreatic cancer, suspected cancer in childhood and young people and other suspected rare cancers. Outcomes of interest are the performance characteristics of the test, particularly sensitivity, specificity and positive and negative predictive values.

Why is this research important

There is very little information currently available on the diagnostic accuracy of tests available in primary care for people with suspected cancer. These studies will inform clinicians on the choice of investigation for symptomatic patients.

2.3 *Cancers insufficiently researched in primary care*

Observational studies of symptomatic primary care patients to estimate the positive predictive value of different symptoms for specific cancers. Priority areas for research are those where the evidence base is currently insufficient and should include prostate cancer, pancreatic cancer, cancer in childhood and young people and other rare cancers. Outcomes of interest are positive predictive values and likelihood ratios for cancer.

Why is this research important

For several cancer sites, the primary care evidence base on the predictive value of symptoms is thin or non-existent. Filling this gap should improve future clinical guidance.

2.4 *Patient experience*

Qualitative studies to assess the key issues in patient experience and patient information needs in the cancer diagnostic pathway, particularly in the interval between first presentation to primary care and first appointment in secondary care. Outcomes of interest are patient satisfaction, quality of life and patient perception of the quality of care and information.

Why is this research important

There was very little information on both patient information needs and patient experience throughout the cancer diagnostic pathway. Filling this gap should improve future patient experience.

3 Other information

3.1 *Scope and how this guideline was developed*

NICE guidelines are developed in accordance with a [scope \[add hyperlink\]](#) that defines what the guideline will and will not cover.

Groups that will be covered

- Children (from birth to 15 years), young adults (aged 16–24 years) and adults (aged 25 years and over) presenting to primary care with signs or symptoms of suspected cancer.

- Subgroups that are identified as needing specific consideration will be considered during development but may include:
 - older people
 - people with cognitive impairment
 - people with multiple morbidities
 - people from lower socioeconomic groups.

Groups that will not be covered

- People who have been referred to secondary care for specialist management.
- People who present for the first time outside of the primary care setting.

How this guideline was developed

NICE commissioned the National Collaborating Centre for Cancer to develop this guideline. The Centre established a Guideline Development Group (see section 4), which reviewed the evidence and developed the recommendations.

The methods and processes for developing NICE clinical guidelines are described in [The guidelines manual](#).

3.2 Related NICE guidance

Details are correct at the time of consultation on the guideline (November 2014). Further information is available on [the NICE website](#).

Published

General

- [Patient experience in adult NHS services](#). NICE clinical guidance 138 (2012).

Condition-specific

- Prostate cancer (update). NICE clinical guideline 175 (2014)

- Familial breast cancer (update). NICE clinical guideline 164 (2013)
- Ovarian cancer. NICE quality standard (2012)
- Lung cancer for adults. NICE quality standard (2012)
- Neutropenic sepsis.. NICE clinical guideline 151 (2012)
- Breast cancer. NICE quality standard (2011)
- [Colorectal cancer](#). NICE clinical guideline 131 (2011).
- [Colonoscopic surveillance for prevention of colorectal cancer in people with ulcerative colitis, Crohn's disease or adenomas](#). NICE clinical guideline 118 (2011).
- [Metastatic malignant disease of unknown primary origin](#). NICE clinical guideline 104 (2010).
- [Lower urinary tract symptoms](#). NICE clinical guideline 97 (2010).
- [Improving outcomes for people with skin tumours including melanoma](#). NICE cancer service guidance (2010).
- [Advanced breast cancer](#). NICE clinical guideline 81 (2009).
- [Early and locally advanced breast cancer](#). NICE clinical guideline 80 (2009).
- [Metastatic spinal cord compression](#). NICE clinical guideline 75 (2008).
- [Improving outcomes for people with skin tumours including melanoma](#). NICE cancer service guidance (2006).
- [Improving outcomes for people with brain and other CNS tumours](#). NICE cancer service guidance (2006).
- [Improving outcomes for people with sarcoma](#). NICE cancer service guidance (2006).
- [Improving outcomes in children and young people with cancer](#). NICE cancer service guidance (2005).
- [Improving supportive and palliative care for adults with cancer](#). NICE cancer service guidance (2004).
- [Improving outcomes in head and neck cancers](#). NICE cancer service guidance (2004).
- [Improving outcomes in colorectal cancer](#). NICE cancer service guidance (2004).

- [Improving outcomes in haematological cancers](#). NICE cancer service guidance (2003).
- [Improving outcomes in urological cancers](#). NICE cancer service guidance (2002).
- [Improving outcomes in breast cancer](#). NICE cancer service guidance (2002).
- Guidance on commissioning cancer services: improving outcomes in lung cancer: the manual. Department of Health (1998). Available from: www.dh.gov.uk
- Improving outcomes in gynaecological cancers. Cancer service guidance (1999). Department of Health, National Cancer Guidance Steering Group. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4005385
- Improving outcomes in upper gastro-intestinal cancers. Cancer service guidance (2001). Department of Health. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4010025

Under development

NICE is developing the following guidance (details available from [the NICE website](#)):

- Bladder cancer. NICE clinical guideline. Publication expected February 2015

The Guideline Development Group, National Collaborating Centre and NICE project team

3.3 *Guideline Development Group*

The Guideline Development Group members listed are those for the 2015 update. For the composition of (the) previous Guideline Development Group(s), see the full guideline.

Lay member ¹¹

Patient/carer member

Nicki Doherty

Lead Cancer Manager Rotherham NHS Foundation Trust¹², General Manager, Barnsley NHS Foundation Trust¹³

Jeanne Fay

General Practitioner, Oxford

Steve Hajioff¹⁴

GDG Chair, Consultant in Public Health Medicine, London

Willie Hamilton

GDG Clinical Lead, Professor of Primary Care Diagnostics, University of Exeter

Susan Hay

Patient and carer member

Georgios (Yoryos) Lyratzopoulos

Senior Clinical Research Associate / Honorary Consultant in Public Health, Department of Public Health and Primary Care, University of Cambridge

David Martin

¹¹ June 2012 – March 2013

¹² January 2012 – June 2013;

¹³ November 2013 - present

¹⁴ September 2013 - present

Patient and carer member

Joan Meakins

General Practitioner, York

Orest Mulka¹⁵

GDG Chair, Retired General Practitioner. Derbyshire

Richard Osborne¹⁶

Consultant Medical Oncologist, Dorset Cancer Centre

Euan Paterson

General Practitioner, Glasgow

Liliana Risi

General Practitioner, London

Karen Sennett

General Practitioner, London

Lindsay Smith

General Practitioner, Somerset

Stuart Williams

Consultant Radiologist, Norfolk & Norwich University Hospital

**3.4 *National Collaborating Centre/ Clinical Guideline
Centre for Cancer***

John Graham

Director

Andrew Champion

Centre Manager

Angela Bennett

¹⁵ January 2012 – June 2013

¹⁶ Chaired meeting 23 & 24 July 2013

Assistant Centre Manager

Katrina Asquith-Coe

Project Manager

Nathan Bromham

Senior Researcher

Mia Schmidt-Hansen

Researcher

Susan O'Connell

Researcher

Laura Bunting

Researcher

David Jarrom

Researcher

Sabine Berendse

Information Specialist

Delyth Morris

Information Specialist

Matthew Prettyjohns

Senior Health Economist

Victoria Kelly

Health Economist

3.5 *NICE project team*

To be completed by NICE

If members of the NICE project team change during guideline development, give details of all people who have fulfilled a role unless told otherwise – see for example CG161.

[Name; style = NICE normal single spacing + bold]

Guideline Lead

[Name; style = NICE normal single spacing + bold]

Clinical Adviser

[Name; style = NICE normal single spacing + bold]

Guideline Commissioning Manager

[Name; style = NICE normal single spacing + bold]

Guideline Coordinator

[Name; style = NICE normal single spacing + bold]

Technical Lead

[Name; style = NICE normal single spacing + bold]

Health Economist

[Name; style = NICE normal single spacing + bold]

Editor

Appendix A: Recommendations from NICE clinical guideline 27 (2005) that have been deleted or changed

Recommendations to be deleted

The table shows recommendations from 2005 that NICE proposed deleting in the 2015 update. The right-hand column gives the replacement recommendation, or explains the reason for the deletion if there is no replacement recommendation.

Recommendation in 2005 guideline	Comment
Patients should be able to consult a primary healthcare professional of the same sex if preferred. [1.1.1]	Recommendation has been deleted because it is not specific to the scope of the guideline
Consideration should be given by the primary healthcare professional to meeting the information and support needs of parents and carers. Consideration should also be given to meeting these particular needs for the people for whom they care, such as children and young people, and people with special needs (for instance, people with learning disabilities or sensory impairment). [1.1.8]	Recommendation has been deleted because this information is already covered by other recommendations
The primary healthcare professional should be aware that some patients find being referred for suspected cancer particularly difficult because of their personal circumstances, such as age, family or work responsibilities, isolation, or other health or social issues. [1.1.9]	Recommendation has been deleted because it is not specific to patient information needs
The primary healthcare professional should be aware that men may have similar support needs to women but may be more reticent about using support services. [1.1.11]	Recommendation has been deleted because it is not specific to the scope of the guideline
In situations where diagnosis or referral has been delayed, or there is significant compromise of the doctor/patient relationship, the primary healthcare professional should take care to assess the information and support needs of the patient, parents and carers, and make sure these needs are met. The patient should be given the opportunity to consult another	Recommendation has been deleted because it is not specific to the scope of the guideline

primary healthcare professional if they wish. [1.1.14]

Primary healthcare professionals should promote awareness of key presenting features of cancer when appropriate. [1.1.15]

Diagnosis of any cancer on clinical grounds alone can be difficult.

Primary healthcare professionals should be familiar with the typical presenting features of cancers, and be able to readily identify these features when patients consult with them. [1.2.1]

Cancers usually present with symptoms commonly associated with benign conditions. The primary healthcare professional should be ready to review the initial diagnosis in patients in whom common symptoms do not resolve as expected. [1.2.2]

Primary healthcare professionals must be alert to the possibility of cancer when confronted by unusual symptom patterns or when patients thought not to have cancer fail to recover as expected. In such circumstances, the primary healthcare professional should systematically review the patient's history and examination, and refer urgently if cancer is a possibility. [1.2.3]

Cancer is uncommon in children, and its detection can present particular difficulties. Primary healthcare professionals should recognise that parents are usually the best observers of their children, and should listen carefully to their concerns. Primary healthcare professionals should also be willing to reassess the initial diagnosis or to seek a second opinion from a colleague if a child fails to recover as expected. [1.2.4]

A patient who presents with symptoms suggestive of cancer should be referred by the primary healthcare professional to a team specialising in the management of the particular type of cancer, depending on local arrangements. [1.2.13]

In patients with features typical of cancer, investigations in primary care should not be allowed to

Recommendation has been deleted as meaning is unclear

Recommendation has been deleted as this was considered to be standard medical practice

Replaced by:
[R10]

Consider a review for people with any symptom that is associated with an increased risk of cancer, but who do not meet the criteria for referral. The review may be:

- planned within a time frame agreed with the person or patient-initiated if their symptoms recur, persist or worsen, new symptoms develop or the person continues to be concerned. [new 2015] [R10]

Replaced by:
[R10]

Consider a review for people with any symptom that is associated with an increased risk of cancer, but who do not meet the criteria for referral. The review may be:

- planned within a time frame agreed with the person or patient-initiated if their symptoms recur, persist or worsen, new symptoms develop or the person continues to be concerned. [new 2015] [R10]

Replaced by:
[R121] & [R122]

Take into account the insight and knowledge of parents and carers when considering urgent referral for a child or young person. [2015] [R121]

Consider referral for children if their parent or carer has persistent concern or anxiety about the child's symptoms, even if the symptoms are most likely to have a benign cause. [2015] [R122]

Replaced by:

Recommendations for each site specific cancer stating where the referral should be made

Replaced by:

Recommendations for each site specific cancer stating what investigations should be performed

delay referral. In patients with less typical symptoms and signs that might, nevertheless, be due to cancer, investigations may be necessary, but should be undertaken urgently to avoid delay. If specific investigations are not readily available locally, an urgent specialist referral should be made. [1.2.14]

A patient who presents with symptoms suggestive of lung cancer should be referred to a team specialising in the management of lung cancer, depending on local arrangements. [1.3.1]

An urgent referral for a chest X-ray should be made when a patient presents with:

- haemoptysis, or
- any of the following unexplained persistent (that is, lasting more than 3 weeks) symptoms and signs:
 - chest and/or shoulder pain
 - dyspnoea
 - weight loss
 - chest signs
 - hoarseness
 - finger clubbing
 - cervical and/or supraclavicular lymphadenopathy
 - cough with or without any of the above
 - features suggestive of metastasis from a lung cancer (for example, in brain, bone, liver or skin).

A report should be made back to the referring primary healthcare professional within 5 days of referral. [1.3.2]

An urgent referral should be made for either of the following:

- persistent haemoptysis in smokers or ex-smokers who are aged 40 years and older
- a chest X-ray suggestive of lung cancer (including pleural effusion and slowly resolving consolidation). [1.3.3]

Immediate referral should be considered for the following:

- signs of superior vena caval obstruction (swelling of the face and/or neck with fixed elevation of jugular venous pressure)
- stridor. [1.3.4]

Replaced by:

[R19] – [R24]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for lung cancer if they:

- have chest X-ray findings that suggest lung cancer or
- are aged over 55 with haemoptysis or
- are aged 40–55, smoke or have smoked in the past, and have haemoptysis or
- are aged 40–55, have never smoked and have haemoptysis and at least one of the following symptoms:
 - cough
 - fatigue
 - shortness of breath
 - chest pain
 - weight loss
 - appetite loss. [new 2015] [R19]

Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who smoke or have smoked in the past and have any of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. [new 2015] [R20]

Offer a full blood count and chest X-ray to assess for lung cancer in people aged 40 and over who have never smoked and have any two of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss
- appetite loss. [new 2015] [R21]

Offer a full blood count to assess for lung cancer in people aged 40 and over who have never smoked and have any of the following unexplained symptoms:

- cough
- fatigue
- shortness of breath
- chest pain
- weight loss

Patients in the following categories have a higher risk of developing lung cancer:

- are current or ex-smokers
- have smoking-related chronic obstructive pulmonary disease (COPD)
- have been exposed to asbestos
- have had a previous history of cancer (especially head and neck).

An urgent referral for a chest X-ray or to a team specialising in the management of lung cancer should be made as for other patients but may be considered sooner, for example if symptoms or signs have lasted for less than 3 weeks. [1.3.5]

Unexplained changes in existing symptoms in patients with underlying chronic respiratory problems should prompt an urgent referral for chest X-ray. [1.3.6]

If the chest X-ray is normal, but there is a high suspicion of lung cancer, patients should be offered an urgent referral. [1.3.7]

In individuals with a history of asbestos exposure and recent onset of chest pain, shortness of breath or unexplained systemic symptoms, lung cancer should be considered and a chest X-ray arranged. If this indicates a pleural effusion, pleural mass or any suspicious lung pathology, an urgent referral should be made. [1.3.8]

A patient who presents with symptoms suggestive of upper gastrointestinal cancer should be referred to a team specialising in the management of upper gastrointestinal cancer, depending on local arrangements. [1.4.1]

An urgent referral for endoscopy or to a specialist with expertise in upper gastrointestinal cancer should be made for patients of any age with dyspepsia³ who present with any of the following:

- chronic gastrointestinal bleeding
- dysphagia
- progressive unintentional weight loss
- persistent vomiting
- iron deficiency anaemia
- epigastric mass

- appetite loss. **[new 2015]** [R22]

Consider an urgent full blood count and chest X-ray (within 2 weeks) to assess for lung cancer in people aged 40 and over with any of the following:

- finger clubbing or
- supraclavicular lymphadenopathy or persistent cervical lymphadenopathy or
- chest signs compatible with lung cancer. **[new 2015]** [R23]

Offer an urgent chest X-ray (within 2 weeks) to assess for lung cancer in people with either:

- thrombocytosis or
- persistent or recurrent chest infection. **[new 2015]** [R24]

Replaced by:
[R29] – [R43]

Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for oesophageal cancer in people:

- aged 45 and over with dysphagia or
- aged 55 and over with weight loss and upper abdominal pain or reflux or dyspepsia. **[new 2015]** [R29]

Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people with haematemesis. **[new 2015]** [R30]

Consider direct access upper gastrointestinal endoscopy to assess for oesophageal cancer in people aged 55 or over with:

- weight loss and nausea or vomiting or
- reflux and nausea or vomiting or
- upper abdominal pain and raised platelet count. **[new]**

- suspicious barium meal result. [1.4.2]

In patients aged 55 years and older with unexplained⁴ and persistent recent-onset dyspepsia alone, an urgent referral for endoscopy should be made. [1.4.3]

In patients aged less than 55 years, endoscopic investigation of dyspepsia is not necessary in the absence of alarm symptoms. [1.4.4]

In patients presenting with dysphagia (interference with the swallowing mechanism that occurs within 5 seconds of having commenced the swallowing process), an urgent referral should be made. [1.4.5]

Helicobacter pylori status should not affect the decision to refer for suspected cancer. [1.4.6]

In patients without dyspepsia, but with unexplained weight loss or iron deficiency anaemia, the possibility of upper gastrointestinal cancer should be recognised and an urgent referral for further investigation considered. [1.4.7]

In patients with persistent vomiting and weight loss in the absence of dyspepsia, upper gastro-oesophageal cancer should be considered and, if appropriate, an urgent referral should be made. [1.4.8]

An urgent referral should be made for patients presenting with either:

- unexplained upper abdominal pain and weight loss, with or without back pain, or
- an upper abdominal mass without dyspepsia. [1.4.9]

In patients with obstructive jaundice an urgent referral should be made, depending on the patient's clinical state. An urgent ultrasound investigation may be considered if available. [1.4.10]

In patients with unexplained worsening of their dyspepsia, an urgent referral should be considered if they have any of the following known risk factors:

- Barrett's oesophagus
- known dysplasia, atrophic gastritis or intestinal metaplasia
- peptic ulcer surgery more than 20 years ago. [1.4.11]

Patients being referred urgently for

2015] [R31]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for pancreatic cancer if they are aged 40 and over and have jaundice.

[new 2015] [R32]

Consider an urgent direct access CT scan (within 2 weeks), or an urgent ultrasound scan if CT is not available, to assess for pancreatic cancer in people aged 60 and over with weight loss and any of the following symptoms:

- diarrhoea
- back pain
- abdominal pain
- nausea or vomiting
- constipation
- new-onset diabetes. **[new 2015] [R33]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with an upper abdominal mass consistent with stomach cancer. **[new 2015] [R36]**

Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with dysphagia. **[new 2015] [R37]**

Offer urgent direct access upper gastrointestinal endoscopy (within 2 weeks) to assess for stomach cancer in people with weight loss who:

- are aged 40 and over, with upper abdominal pain lasting 2 weeks or more, and nausea or vomiting or
- are aged 55 and over with upper abdominal pain, reflux or dyspepsia. **[new 2015] [R38]**

Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people with weight loss who:

- also have appetite loss or
- are aged under 55 with dyspepsia or upper abdominal pain lasting 2 weeks or more or
- are aged 55 and over with nausea or vomiting. **[new 2015] [R39]**

Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with reflux and nausea or vomiting. **[new 2015] [R40]**

Consider direct access upper gastrointestinal endoscopy to assess for stomach cancer in people aged 55 and over with upper abdominal pain and raised platelet counts. **[new 2015] [R41]**

Consider urgent direct access ultrasound (within 2 weeks) to assess for gall bladder cancer in people with an upper abdominal mass consistent with an enlarged gall bladder. **[new 2015] [R42]**

endoscopy should ideally be free from acid suppression medication, including proton pump inhibitors or H2 receptor antagonists, for a minimum of 2 weeks. [1.4.12]

In patients where the decision to refer has been made, a full blood count may assist specialist assessment in the outpatient clinic.

This should be carried out in accordance with local arrangements. [1.4.13]

All patients with new-onset dyspepsia should be considered for a full blood count in order to detect iron deficiency anaemia. [1.4.14]

A patient who presents with symptoms suggestive of colorectal or anal cancer should be referred to a team specialising in the management of lower gastrointestinal cancer, depending on local arrangements. [1.5.1]

In patients with equivocal symptoms who are not unduly anxious, it is reasonable to use a period of 'treat, watch and wait' as a method of management. [1.5.2]
In patients aged 40 years and older, reporting rectal bleeding with a change of bowel habit towards looser stools and/or increased stool frequency persisting for 6 weeks or more, an urgent referral should be made. [1.5.3]

In patients aged 60 years and older, with rectal bleeding persisting for 6 weeks or more without a change in bowel habit and without anal symptoms, an urgent referral should be made. [1.5.5.]

In patients aged 60 years and older, with a change in bowel habit to looser stools and/or more frequent stools persisting for 6 weeks or more without rectal bleeding, an urgent referral should be made. [1.5.6]

In patients presenting with a right lower abdominal mass consistent with involvement of the large bowel, an urgent referral should be made, irrespective of age. [1.5.7]

In patients presenting with a palpable rectal mass (intraluminal and not pelvic), an urgent referral should be made, irrespective of age. (A pelvic mass outside the

Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for liver cancer in people with an upper abdominal mass consistent with an enlarged liver. **[new 2015]** [R43]

Replaced by:
[R44] – [R51]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 50 and have unexplained rectal bleeding. **[new 2015]** [R44]

Refer people using a suspected cancer pathway referral (with an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women). **[new 2015]** [R45]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 60 and have unexplained changes in their bowel habit. **[new 2015]** [R46]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people with a rectal or abdominal mass. **[new 2015]** [R47]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if they are aged over 40 with unexplained weight loss and abdominal pain. **[new 2015]** [R48]

Offer testing for occult blood in faeces to assess for colorectal cancer in people without rectal bleeding who:

- have abdominal pain or
- have weight loss or
- are aged under 60 and have a change in bowel habit or iron-deficiency anaemia (with haemoglobin levels of 12 g/dl or below for men and 11 g/dl or below for women). **[new 2015]** [R49]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer if tests show occult blood in their faeces. **[new**

bowel would warrant an urgent referral to a urologist or gynaecologist.) [1.5.8]
In men of any age with unexplained iron deficiency anaemia and a haemoglobin of 11 g/100 ml or below, an urgent referral should be made. [1.5.9]
In non-menstruating women with unexplained iron deficiency anaemia and a haemoglobin of 10 g/100 ml or below, an urgent referral should be made. [1.5.10]
In patients with ulcerative colitis or a history of ulcerative colitis, a plan for follow-up should be agreed with a specialist and offered to the patient as a normal procedure in an effort to detect colorectal cancer in this high-risk group. [1.5.11]

There is insufficient evidence to suggest that a positive family history of colorectal cancer can be used as a criterion to assist in the decision about referral of a symptomatic patient. [1.5.12]

In patients with equivocal symptoms, a full blood count may help in identifying the possibility of colorectal cancer by demonstrating iron deficiency anaemia, which should then determine if a referral should be made and its urgency. [1.5.13]

In patients for whom the decision to refer has been made, a full blood count may assist specialist assessment in the outpatient clinic. This should be in accordance with local arrangements. [1.5.14]

In patients for whom the decision to refer has been made, no examinations or investigations other than those referred to earlier (abdominal and rectal examination, full blood count) are recommended as this may delay referral. [1.5.15]

A patient who presents with symptoms suggestive of breast cancer should be referred to a team specialising in the management of breast cancer. [1.6.1]

In most cases, the definitive diagnosis will not be known at the time of referral, and many patients who are referred will be found not to have cancer. However, primary healthcare professionals should

2015] [R50]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for colorectal cancer in people aged under 50 with rectal bleeding and any of the following unexplained symptoms or findings:

- abdominal pain
- change in bowel habit
- weight loss
- iron-deficiency anaemia (haemoglobin levels 12 g/dl or below for men and 11 g/dl or below for women).

[new 2015] [R51]

Replaced by:

[R53] – [R54]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 30 and over and have a breast lump with or without pain. **[new 2015]** [R53]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for breast cancer if they are aged 50 and over with any of the following symptoms in one nipple only:

convey optimism about the effectiveness of treatment and survival because a patient being referred with a breast lump will be naturally concerned. [1.6.2]

People of all ages who suspect they have breast cancer may have particular information and support needs. The primary healthcare professional should discuss these needs with the patient and respond sensitively to them. [1.6.3]

Primary healthcare professionals should encourage all patients, including women over 50 years old, to be breast aware in order to minimise delay in the presentation of symptoms.[1.6.4]

A woman's first suspicion that she may have breast cancer is often when she finds a lump in her breast. The primary healthcare professional should examine the lump with the patient's consent.

The features of a lump that should make the primary healthcare professional strongly suspect cancer are a discrete, hard lump with fixation, with or without skin tethering. In patients presenting in this way an urgent referral should be made, irrespective of age.

[1.6.5]

In a woman aged 30 years and older with a discrete lump that persists after her next period, or presents after menopause, an urgent referral should be made.

[1.6.6]

Breast cancer in women aged younger than 30 years is rare, but does occur. Benign lumps (for example, fibroadenoma) are common, however, and a policy of referring these women urgently would not be appropriate; instead, non-urgent referral should be considered. However, in women aged younger than 30 years:

- with a lump that enlarges, or
- with a lump that has other features associated with cancer (fixed and hard), or
- in whom there are other reasons for concern such as family history

an urgent referral should be made.

[1.6.7]

The patient's history should always be taken into account. For

- discharge
- retraction
- other changes of concern. [new 2015] [R54]

example, it may be appropriate, in discussion with a specialist, to agree referral within a few days in patients reporting a lump or other symptom that has been present for several months. [1.6.8]

In a patient who has previously had histologically confirmed breast cancer, who presents with a further lump or suspicious symptoms, an urgent referral should be made, irrespective of age. [1.6.9]

In patients presenting with unilateral eczematous skin or nipple change that does not respond to topical treatment, or with nipple distortion of recent onset, an urgent referral should be made. [1.6.10]

In patients presenting with spontaneous unilateral bloody nipple discharge, an urgent referral should be made. [1.6.11]

Breast cancer in men is rare and is particularly rare in men under 50 years of age. However, in a man aged 50 years and older with a unilateral, firm subareolar mass with or without nipple distortion or associated skin changes, an urgent referral should be made. [1.6.12]

In patients presenting with symptoms and/or signs suggestive of breast cancer, investigation prior to referral is not recommended. [1.6.13]

In patients presenting solely with breast pain, with no palpable abnormality, there is no evidence to support the use of mammography as a discriminatory investigation for breast cancer.

Therefore, its use in this group of patients is not recommended. Non-urgent referral may be considered in the event of failure of initial treatment and/or unexplained persistent symptoms. [1.6.14]

A patient who presents with symptoms suggesting gynaecological cancer should be referred to a team specialising in the management of gynaecological cancer, depending on local arrangements. [1.7.1]

Replaced by:
[R56] – [R69] & [R71]

Refer the woman urgently¹⁷ if physical examination identifies ascites and/or a pelvic or abdominal mass (which is not obviously uterine fibroids). [R56]

¹⁷ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

The first symptoms of gynaecological cancer may be alterations in the menstrual cycle, intermenstrual bleeding, postcoital bleeding, postmenopausal bleeding or vaginal discharge.

When a patient presents with any of these symptoms, the primary healthcare professional should undertake a full pelvic examination, including speculum examination of the cervix. [1.7.2] In patients found on examination of the cervix to have clinical features that raise the suspicion of cervical cancer, an urgent referral should be made. A cervical smear test is not required before referral, and a previous negative cervical smear result is not a reason to delay referral. [1.7.3]

This recommendation has been updated and replaced by section 1.1. in 'Ovarian cancer: the diagnosis and initial management of ovarian cancer' (NICE clinical guideline 122, 2011) [1.7.4]

Any woman with a palpable abdominal or pelvic mass on examination that is not obviously uterine fibroids or not of gastrointestinal or urological origin should have an urgent ultrasound scan. If the scan is suggestive of cancer, or if ultrasound is not available, an urgent referral should be made. [1.7.5]

When a woman who is not on hormone replacement therapy presents with postmenopausal bleeding, an urgent referral should be made. [1.7.6]

When a woman on hormone replacement therapy presents with persistent or unexplained postmenopausal bleeding after cessation of hormone replacement therapy for 6 weeks, an urgent referral should be made. [1.7.7.]

Tamoxifen can increase the risk of endometrial cancer. When a woman taking tamoxifen presents with postmenopausal bleeding, an urgent referral should be made. [1.7.8]

Carry out tests in primary care if a woman (especially if 50 or over) reports having any of the following symptoms on a persistent or frequent basis – particularly more than 12 times per month:

- persistent abdominal distension (women often refer to this as 'bloating')
- feeling full (early satiety) and/or loss of appetite
- pelvic or abdominal pain
- increased urinary urgency and/or frequency. [R57]

Consider carrying out tests in primary care if a woman reports unexplained weight loss, fatigue or changes in bowel habit. [R58]

Advise any woman who is not suspected of having ovarian cancer to return to her GP if her symptoms become more frequent and/or persistent. [R59]

Carry out appropriate tests for ovarian cancer in any woman of 50 or over who has experienced symptoms within the last 12 months that suggest irritable bowel syndrome (IBS)¹⁸, because IBS rarely presents for the first time in women of this age. [R60]

Measure serum CA125 in primary care in women with symptoms that suggest ovarian cancer. [R61]

If serum CA125 is 35 IU/ml or greater, arrange an ultrasound scan of the abdomen and pelvis. [R62]

If the ultrasound suggests ovarian cancer, refer the woman urgently¹⁹ for further investigation. [R63]

For any woman who has normal serum CA125 (less than 35 IU/ml), or CA125 of 35 IU/ml or greater but a normal ultrasound:

- assess her carefully for other clinical causes of her symptoms and investigate if appropriate
- if no other clinical cause is apparent, advise her to return to her GP if her symptoms become more frequent and/or persistent. [R64]

Refer women using a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer if they are aged 55 and over with post-menopausal bleeding (unexplained vaginal bleeding more than 12 months after menstruation has stopped because of the menopause). [new 2015] [R65]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for endometrial cancer in women aged under 55 with post-menopausal bleeding. [new 2015] [R66]

¹⁸ See Irritable bowel syndrome in adults (NICE clinical guideline 61).

¹⁹ An urgent referral means that the woman is referred to a gynaecological cancer service within the national target in England and Wales for referral for suspected cancer, which is currently 2 weeks.

An urgent referral should be considered in a patient with persistent intermenstrual bleeding and a negative pelvic examination. [1.7.9]

Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with unexplained symptoms of vaginal discharge who:

- are presenting with these symptoms for the first time or
- have a high platelet count or
- report haematuria. **[new 2015]** **[R67]**

Consider direct access ultrasound scan to assess for endometrial cancer in women aged 55 and over with visible haematuria and any of the following:

- low haemoglobin levels or
- high platelet count or
- high blood glucose levels. **[new 2015]** **[R68]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for women if the appearance of their cervix is consistent with cervical cancer. **[new 2015]** **[R69]**

Consider a suspected cancer pathway referral (with an appointment within 2 weeks) for vaginal cancer in women with an unexplained palpable mass in or at the entrance to the vagina. **[new 2015]** **[R71]**

Replaced by:
[R70]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for vulval cancer in women with an unexplained vulval lump, ulceration or bleeding. **[new 2015]**

When a woman presents with vulval symptoms, a vulval examination should be offered. If an unexplained vulval lump is found, an urgent referral should be made. [1.7.10]

Vulval cancer can also present with vulval bleeding due to ulceration. A patient with these features should be referred urgently. [1.7.11]

Vulval cancer may also present with pruritus or pain. For a patient who presents with these symptoms, it is reasonable to use a period of 'treat, watch and wait' as a method of management. But this should include active follow-up until symptoms resolve or a diagnosis is confirmed. If symptoms persist, the referral may be urgent or non-urgent, depending on the symptoms and the degree of concern about cancer. [1.7.12]

A patient who presents with symptoms or signs suggestive of urological cancer should be referred to a team specialising in the management of urological cancer, depending on local arrangements. [1.8.1]

Patients presenting with symptoms suggesting prostate cancer should have a digital rectal examination (DRE) and prostate-specific

Replaced by:
[R72] – [R74]

Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for prostate cancer if their prostate feels malignant on digital rectal examination. **[new 2015]** **[R72]**

Consider a prostate-specific antigen (PSA) test and digital rectal examination to assess for prostate cancer

antigen (PSA) test after counselling. Symptoms will be related to the lower urinary tract and may be inflammatory or obstructive. [1.8.2]

Prostate cancer is also a possibility in male patients with any of the following unexplained symptoms:

- erectile dysfunction
- haematuria
- lower back pain
- bone pain
- weight loss, especially in the elderly.

These patients should also be offered a DRE and a PSA test.

[1.8.3]

Urinary infection should be excluded before PSA testing, especially in men presenting with lower tract symptoms. The PSA test should be postponed for at least 1 month after treatment of a proven urinary infection. [1.8.4]

If a hard, irregular prostate typical of a prostate carcinoma is felt on rectal examination, then the patient should be referred urgently. The PSA should be measured and the result should accompany the referral. Patients do not need urgent referral if the prostate is simply enlarged and the PSA is in the age-specific reference range.

[1.8.5]

In a male patient with or without lower urinary tract symptoms and in whom the prostate is normal on DRE but the age-specific PSA is raised or rising, an urgent referral should be made. In those patients whose clinical state is compromised by other comorbidities, a discussion with the patient or carers and/or a specialist in urological cancer may be more appropriate. [1.8.6]

Symptomatic patients with high PSA levels should be referred urgently. [1.8.7]

If there is doubt about whether to refer an asymptomatic male with a borderline level of PSA, the PSA test should be repeated after an interval of 1 to 3 months. If the second test indicates that the PSA level is rising, the patient should be referred urgently. [1.8.9]

Male or female adult patients of any age who present with painless

in men with:

- any lower urinary tract symptoms, such as nocturia, urinary frequency, hesitancy, urgency or retention or
- erectile dysfunction or
- visible haematuria. **[new 2015]** [R73]

Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for prostate cancer if their PSA levels are above the age-specific reference range. **[new 2015]** [R74]

Replaced by:
[R75] – [R80]

macroscopic haematuria should be referred urgently. [1.8.9]

In male or female patients with symptoms suggestive of a urinary infection who also present with macroscopic haematuria,

investigations should be undertaken to diagnose and treat the infection before consideration of referral. If infection is not confirmed the patient should be referred urgently. [1.8.10]

In all adult patients aged 40 years and older who present with recurrent or persistent urinary tract infection associated with haematuria, an urgent referral should be made. [1.8.11]

In patients under 50 years of age with microscopic haematuria, the urine should be tested for proteinuria and serum creatinine levels measured. Those with proteinuria or raised serum creatinine should be referred to a renal physician. If there is no proteinuria and serum creatinine is normal, a non-urgent referral to a urologist should be made. [1.8.12]

In patients aged 50 years and older who are found to have unexplained microscopic haematuria, an urgent referral should be made. [1.8.13]

Any patient with an abdominal mass identified clinically or on imaging that is thought to be arising from the urinary tract should be referred urgently.

[1.8.14]

Any patient with a swelling or mass in the body of the testis should be referred urgently. [1.8.15]

An urgent ultrasound should be considered in men with a scrotal mass that does not transilluminate and/or when the body of the testis cannot be distinguished. [1.8.16]

An urgent referral should be made for any patient presenting with symptoms or signs of penile cancer. These include progressive ulceration or a mass in the glans or prepuce particularly, but can involve the skin of the penile shaft. Lumps within the corpora

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have unexplained visible haematuria without urinary tract infection. **[new 2015]**

[R75]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. **[new 2015]**

[R76]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for bladder cancer if they are aged 60 and over and have unexplained non-visible haematuria and either dysuria or a raised white cell count on a blood test. **[new 2015]**

[R77]

Consider referral for bladder cancer in people aged 60 and over with recurrent or persistent urinary tract infection that is unexplained. **[new 2015]**

[R78]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have unexplained visible haematuria. **[new 2015]**

[R79]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for renal cancer if they are aged 45 and over and have visible haematuria that persists or recurs after successful treatment of urinary tract infection. **[new 2015]**

[R80]

Replaced by:

[R85] – [R86]

Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for testicular cancer if they have a non-painful enlargement or change in shape or texture of the testis. **[new 2015]**

[R85]

Consider a direct access ultrasound scan as part of clinical reassessment for testicular cancer in men with unexplained or persistent testicular symptoms. **[new 2015]**

[R86]

Replaced by:

[R87] – [R89]

Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a penile mass or ulcerated lesion and sexually transmitted infection has been excluded as a cause. **[new 2015]**

[R87]

cavernosa not involving penile skin are usually not cancer but indicate Peyronie's disease, which does not require urgent referral. [1.8.17]

A patient who presents with symptoms suggesting haematological cancer should be referred to a team specialising in the management of haematological cancer, depending on local arrangements. [1.9.1] Primary healthcare professionals should be aware that haematological cancers can present with a variety of symptoms that may have a number of different clinical explanations. [1.9.2]

Combinations of the following symptoms and signs may suggest haematological cancer and warrant full examination, further investigation (including a blood count and film) and possible referral:

- fatigue
- drenching night sweats
- fever
- weight loss
- generalised itching
- breathlessness
- bruising
- bleeding
- recurrent infections
- bone pain
- alcohol-induced pain
- abdominal pain
- lymphadenopathy
- splenomegaly.

The urgency of referral depends on the severity of the symptoms and signs, and findings of investigations. [1.9.3]

In patients with a blood count or blood film reported as acute leukaemia, an immediate referral should be made. [1.9.4]

In patients with persistent unexplained splenomegaly, an urgent referral should be made. [1.9.5]

Investigation of patients with

Refer men using a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer if they have a persistent penile lesion after treatment for a sexually transmitted infection has been completed.

[new 2015] [R88]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for penile cancer in men with unexplained or persistent symptoms affecting the foreskin or glands. **[new 2015] [R89]**

Replaced by:

[R105] – [R108], [Ra1], [R109] – [R112]

Consider a very urgent full blood count (within 48 hours) to assess for leukaemia in adults with any of the following symptoms:

- pallor
- persistent fatigue
- unexplained fever
- unexplained persistent or recurrent infection
- generalised lymphadenopathy
- unexplained bruising
- unexplained bleeding
- unexplained petechiae
- hepatosplenomegaly. **[new 2015] [R105]**

Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young adults with any of the following symptoms:

- pallor
- persistent fatigue
- unexplained fever
- unexplained persistent infection
- generalised lymphadenopathy
- persistent or unexplained bone pain
- unexplained bruising
- unexplained bleeding. **[new 2015] [R106]**

Refer children and young adults for immediate specialist assessment for leukaemia if they have unexplained petechiae or hepatosplenomegaly. **[new 2015] [R107]**

Offer a full blood count, blood tests for calcium and plasma viscosity or erythrocyte sedimentation rate to assess for myeloma in people aged 60 and over with persistent bone pain, particularly back pain or unexplained fracture. **[new 2015] [R108]**

Offer very urgent protein electrophoresis (within 48 hours) to assess for myeloma in people aged 60 and over with hypercalcaemia or leucopenia and a presentation that is consistent with possible myeloma.

[new 2015] [Ra1]

persistent unexplained fatigue should include a full blood count, blood film and erythrocyte sedimentation rate, plasma viscosity or C-reactive protein (according to local policy), and repeated at least once if the patient's condition remains unexplained and does not improve. [1.9.6]

Investigation of patients with unexplained lymphadenopathy should include a full blood count, blood film and erythrocyte sedimentation rate, plasma viscosity or C-reactive protein (according to local policy). [1.9.7] Any of the following additional features of lymphadenopathy should trigger further investigation and/or referral:

- persistence for 6 weeks or more
- lymph nodes increasing in size
- lymph nodes greater than 2 cm in size
- widespread nature
- associated splenomegaly, night sweats or weight loss. [1.9.8]

Investigation of a patient with unexplained bruising, bleeding, and purpura or symptoms suggesting anaemia should include a full blood count, blood film, clotting screen and erythrocyte sedimentation rate, plasma viscosity or C-reactive protein (according to local policy). [1.9.9]

A patient with bone pain that is persistent and unexplained should be investigated with full blood count and X-ray, urea and electrolytes, liver and bone profile, PSA test (in males) and erythrocyte sedimentation rate, plasma viscosity or C-reactive protein (according to local policy). [1.9.10]

In patients with spinal cord compression or renal failure suspected of being caused by myeloma, an immediate referral should be made. [1.9.11]

A patient presenting with skin lesions suggestive of skin cancer or in whom a biopsy has been confirmed should be referred to a team specialising in skin cancer. [1.10.1]

Consider very urgent protein electrophoresis (within 48 hours) to assess for myeloma if the plasma viscosity or erythrocyte sedimentation rate and presentation are consistent with possible myeloma. **[new 2015]** [R109]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if the results of protein electrophoresis suggest myeloma. **[new 2015]** [R110]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. **[new 2015]** [R111]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. **[new 2015]** [R112]

Replaced by:
[R90], [Ra2], [R92] – [R95]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) if dermatoscopy suggests malignant melanoma of the skin. **[new 2015]**

All primary healthcare professionals should be aware of the 7-point weighted checklist (see recommendation 1.10.8) for assessment of pigmented skin lesions. [1.10.2]

All primary healthcare professionals who perform minor surgery should have received appropriate accredited training in relevant aspects of skin surgery including cryotherapy, curettage, and incisional and excisional biopsy techniques, and should undertake appropriate continuing professional development. [1.10.3] Patients with persistent or slowly evolving unresponsive skin conditions in which the diagnosis is uncertain and cancer is a possibility should be referred to a dermatologist. [1.10.4]

All excised skin specimens should be sent for pathological examination. [1.10.5]

On making a referral of a patient in whom an excised lesion has been diagnosed as malignant, a copy of the pathology report should be sent with the referral

correspondence, as there may be details (such as tumour thickness, excision margin) that will specifically influence future management. [1.10.6]

Change is a key element in diagnosing malignant melanoma.

For low-suspicion lesions, careful monitoring for change should be undertaken using the 7-point checklist (see recommendation 1.10.8) for 8 weeks. Measurement should be made with photographs and a marker scale and/or ruler. [1.10.7]

All primary healthcare professionals should use the weighted 7-point checklist in the assessment of pigmented lesions to determine referral:

Major features of the lesions:

- change in size
- irregular shape
- irregular colour.

Minor features of the lesions:

[R90]

Refer people using a suspected cancer pathway referral (for an appointment within 2 weeks) for malignant melanoma if they present with a suspicious pigmented skin lesion that has a weighted 7-point checklist score of 3 or more.

Major features of the lesions (scoring 2 points each):

- change in size
- irregular shape
- irregular colour.

Minor features of the lesions (scoring 1 point each):

- largest diameter 7 mm or more
- inflammation
- oozing
- change in sensation. [new 2015] [Ra2]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a skin lesion that raises the suspicion of squamous cell carcinoma. [new 2015] [R92]

Consider routine referral for people if they have a skin lesion that raises the suspicion of a basal cell carcinoma²⁰. [new 2015] [R93]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for basal cell carcinoma if there is concern that a delay may have an unfavourable impact, because of factors such as lesion site or size. [new 2015] [R94]

GPs should only excise suspected basal cell carcinomas in accordance with NICE guidance on Improving outcomes for people with skin tumours including melanoma. [new 2015] [R95]

²⁰

Typical features of basal cell carcinoma include: an ulcer with a raised rolled edge; prominent fine blood vessels around a lesion; or a nodule on the skin (particularly pearly or waxy nodules).

- largest diameter 7 mm or more
- inflammation
- oozing
- change in sensation.

Suspicion is greater for lesions scoring 3 points or more (based on major features scoring 2 points each and minor features scoring 1 point each). However, if there are strong concerns about cancer, any one feature is adequate to prompt urgent referral. [1.10.8]

In patients with a lesion suspected to be melanoma (see recommendation 1.10.8), an urgent referral to a dermatologist or other suitable specialist with experience of melanoma diagnosis should be made, and excision in primary care should be avoided [1.10.9]

Squamous cell carcinomas present as keratinizing or crusted tumours that may ulcerate. Non-healing lesions larger than 1 cm with significant induration on palpation, commonly on face, scalp or back of hand with a documented expansion over 8 weeks, may be squamous cell carcinomas and an urgent referral should be made. [1.10.10]

Squamous cell carcinomas are common in patients on immunosuppressive treatment, but may be atypical and aggressive. In patients who have had an organ transplant who develop new or growing cutaneous lesions, an urgent referral should be made. [1.10.11]

In any patient with histological diagnosis of a squamous cell carcinoma made in primary care, an urgent referral should be made. [1.10.12]

Basal cell carcinomas are slow growing, usually without significant expansion over 2 months, and usually occur on the face. Where there is a suspicion that the patient has a basal cell carcinoma, a non-urgent referral should be made. [1.10.13]

All pigmented lesions that are not viewed as suspicious of melanoma but are excised should have a lateral excision margin of 2 mm of clinically normal skin and cut to include subcutaneous fat in depth.

[1.10.14]

A patient who presents with symptoms suggestive of head and neck or thyroid cancer should be referred to an appropriate specialist or the neck lump clinic, depending on local arrangements.

[1.11.1]

Any patient with persistent symptoms or signs related to the oral cavity in whom a definitive diagnosis of a benign lesion cannot be made should be referred or followed up until the symptoms and signs disappear. If the symptoms and signs have not disappeared after 6 weeks, an urgent referral should be made.

[1.11.2]

Primary healthcare professionals should advise all patients, including those with dentures, to have regular dental checkups.

[1.11.3]

In a patient who presents with unexplained red and white patches (including suspected lichen planus) of the oral mucosa that are:

- painful, or
- swollen, or
- bleeding

an urgent referral should be made.

A non-urgent referral should be made in the absence of these features. If oral lichen planus is confirmed, the patient should be monitored for oral cancer as part of routine dental examination.

[1.11.4]

In patients with unexplained ulceration of the oral mucosa or mass persisting for more than 3 weeks, an urgent referral should be made. [1.11.5]

In adult patients with unexplained tooth mobility persisting for more than 3 weeks, an urgent referral to a dentist should be made. [1.11.6]

In any patient with hoarseness persisting for more than 3 weeks, particularly smokers aged 50 years and older and heavy drinkers, an urgent referral for a chest X-ray should be made. Patients with positive findings should be referred urgently to a team specialising in the management of lung cancer. Patients with a negative finding should be urgently referred to a team specialising in head and neck

Replaced by:

[R96] – [R101]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with persistent unexplained hoarseness. **[new 2015] [R96]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for laryngeal cancer in people aged 45 and over with an unexplained lump in the neck. **[new 2015] [R97]**

Consider an urgent referral (for an appointment within 2 weeks) for assessment for oral cancer by the community dental service in people with an unexplained lump on the lip or in the oral cavity that has not been assessed by a dental surgeon. **[new 2015] [R98]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people with a lump on the lip or in the oral cavity that has been assessed by a dental surgeon to be consistent with oral cancer. **[new 2015] [R99]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with unexplained ulceration in the oral cavity lasting for more than 14 days. **[new 2015] [R100]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for oral cancer in people with a persistent and unexplained lump in the neck. **[new 2015] [R101]**

cancer. [1.11.7]

In patients with an unexplained lump in the neck which has recently appeared or a lump which has not been diagnosed before that has changed over a period of 3 to 6 weeks, an urgent referral should be made. [1.11.8]

In patients with an unexplained persistent swelling in the parotid or submandibular gland, an urgent referral should be made. [1.11.9]

In patients with unexplained persistent sore or painful throat, an urgent referral should be made.

[1.11.10]

In patients with unilateral unexplained pain in the head and neck area for more than 4 weeks, associated with otalgia (ear ache) but with normal otoscopy, an urgent referral should be made.

[1.11.11]

With the exception of persistent hoarseness (see recommendation 1.11.7), investigations for head and neck cancer in primary care are not recommended as they can delay referral. [1.11.12]

In patients presenting with symptoms of tracheal compression including stridor due to thyroid swelling, immediate referral should be made. [1.11.13]

In patients presenting with a thyroid swelling associated with any of the following, an urgent referral should be made:

- a solitary nodule increasing in size
- a history of neck irradiation
- a family history of an endocrine tumour
- unexplained hoarseness or voice changes
- cervical lymphadenopathy
- very young (pre-pubertal) patients
- patients aged 65 years and older. [1.11.14]

In patients with a thyroid swelling without stridor or any of the features indicated in recommendation 1.11.14, the primary healthcare professional should request thyroid function tests. Patients with hyper- or hypothyroidism and an associated goitre are very unlikely to have thyroid cancer and could be

Replaced by:
[R102]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for thyroid cancer in people with an unexplained thyroid lump. **[new 2015]**
[R102]

referred, nonurgently, to an endocrinologist. Those with goitre and normal thyroid function tests who do not have any of the features indicated in recommendation 1.11.14 should be referred non-urgently. [1.11.15] Initiation of other investigations by the primary healthcare professional, such as ultrasonography or isotope scanning, is likely to result in unnecessary delay and is not recommended. [1.11.16]

A patient who presents with symptoms suggestive of brain or CNS cancer should be referred to an appropriate specialist, depending on local arrangements. [1.12.1]

If a primary healthcare professional has concerns about the interpretation of a patient's symptoms and/or signs, a discussion with a local specialist should be considered. If rapid access to scanning is available, this investigation should also be considered as an alternative. [1.12.2]

In patients with new, unexplained headaches or neurological symptoms, the primary healthcare professional should undertake a neurological examination guided by the symptoms, but including examination for papilloedema. The absence of papilloedema does not exclude the possibility of a brain tumour. [1.12.3]

In any patient with symptoms related to the CNS (including progressive neurological deficit, new-onset seizures, headaches, mental changes, cranial nerve palsy, unilateral sensorineural deafness) in whom a brain tumour is suspected, an urgent referral should be made. The development of new signs related to the CNS should be considered as potential indications for referral. [1.12.4]

In patients with headaches of recent onset accompanied by either features suggestive of raised intracranial pressure (for example, vomiting, drowsiness, posture-related headache, headache with pulse-synchronous tinnitus) or other focal or non-focal

Replaced by:
[R103] – [R104]

Consider an urgent direct access MRI scan of the brain (within 2 weeks) to assess for brain or central nervous system cancer in adults with progressive, sub-acute loss of central neurological function. **[new 2015] [R103]**

Consider very urgent referral (for an appointment within 48 hours) for suspected brain or central nervous system cancer in children and young adults with newly abnormal cerebellar or other central neurological function. **[new 2015] [R104]**

neurological symptoms (for example, blackout, change in personality or memory), an urgent referral should be made. [1.12.5]

In patients with unexplained headaches of recent onset, present for at least 1 month but not accompanied by features suggestive of raised intracranial pressure (see recommendation 1.12.5),

discussion with a local specialist or referral (usually non-urgent) should be considered. [1.12.6]

In patients with a new, qualitatively different unexplained headache that becomes progressively severe, an urgent referral should be made. [1.12.7]

Re-assessment and re-examination is required if the patient does not progress according to expectations. [1.12.8]

A detailed history should be taken from the patient and an eyewitness to the event if possible, to determine whether or not a seizure is likely to have occurred. [1.12.9]

In patients presenting with a seizure, a physical examination (including cardiac, neurological, mental state) and developmental assessment, where appropriate, should be carried out. [1.12.10]

In any patient with suspected recent-onset seizures, an urgent referral to a neurologist should be made. [1.12.11]

In patients with rapid progression of:

- subacute focal neurological deficit
- unexplained cognitive impairment, behavioural disturbance, or slowness or a combination of these
- personality changes confirmed by a witness (for example, a carer, friend or a family member) and for which there is no reasonable explanation even in the absence of the other symptoms and signs of a brain tumour

an urgent referral to an appropriate specialist should be considered. [1.12.12]

In patients previously diagnosed with any cancer an urgent referral should be made if the patient

develops any of the following symptoms:

- recent-onset seizure
- progressive neurological deficit
- persistent headaches
- new mental or cognitive changes
- new neurological signs.

[1.12.13]

A patient who presents with symptoms suggesting bone cancer or sarcoma should be referred to a team specialising in the management of bone cancer and sarcoma, or to a recognised bone cancer centre, depending on local arrangements. [1.13.1]

If a primary healthcare professional has concerns about the interpretation of a patient's symptoms and/or signs, a discussion with the local specialist should be considered. [1.13.2]
Patients with increasing, unexplained or persistent bone pain or tenderness, particularly pain at rest (and especially if not in the joint), or an unexplained limp should be investigated by the primary healthcare professional urgently. The nature of the investigations will vary according to the patient's age and clinical features.

* In older people metastases, myeloma or lymphoma, as well as sarcoma, should be considered. [1.13.3]

A patient with a suspected spontaneous fracture should be referred for an immediate X-ray. [1.13.4]

If an X-ray indicates that bone cancer is a possibility, an urgent referral should be made. [1.13.5]

If the X-ray is normal but symptoms persist, the patient should be followed up and/or a repeat X-ray or bone function tests or a referral requested. [1.13.6]

In patients presenting with a palpable lump, an urgent referral for suspicion of soft tissue sarcoma should be made if the lump is:

- greater than about 5 cm in diameter
- deep to fascia, fixed or

Replaced by:

[R113] – [R116]

Consider an urgent direct access X-ray (within 2 weeks) to assess for bone sarcoma in children and young adults with unexplained bone swelling or pain. **[new 2015] [R113]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if an X-ray suggests the possibility of bone sarcoma. **[new 2015] [R114]**

Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for soft tissue sarcoma in people with an unexplained lump that is increasing in size. **[new 2015] [R115]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if they have ultrasound scan findings that are suggestive of soft tissue sarcoma or if ultrasound findings are uncertain and clinical concern persists. **[new 2015] [R116]**

- immobile
- painful
- increasing in size
- a recurrence after previous excision.

If there is any doubt about the need for referral, discussion with a local specialist should be undertaken. [1.13.7]

If a patient has HIV disease, Kaposi's sarcoma should be considered and a referral made if this is suspected. [1.13.8]

Children and young people who present with symptoms and signs of cancer should be referred to a paediatrician or a specialist children's cancer service, if appropriate. [1.14.1]

Replaced by:
[R117] – [R122]

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]**
[R117]

Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex.
[new 2015] **[R118]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]** **[R119]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. **[new 2015]** **[R120]**

Take into account the insight and knowledge of parents and carers when considering urgent referral for a child or young person. **[new 2015]** **[R121]**

Consider referral for children if their parent or carer has persistent concern or anxiety about the child's symptoms, even if the symptoms are most likely to have a benign cause. **[new 2015]** **[R122]**

Replaced by:
[R117] – [R122]

Childhood cancer is rare and may present initially with symptoms and signs associated with common conditions. Therefore, in the case of a child or young person presenting several times (for example, three or more times) with the same problem, but with no clear diagnosis, urgent referral should be made. [1.14.2]

The parent is usually the best observer of the child's or young person's symptoms. The primary healthcare professional should take note of parental insight and

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]**
[R117]

Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex.
[new 2015] **[R118]**

Consider very urgent referral (for an appointment within

knowledge when considering urgent referral. [1.14.3]

Persistent parental anxiety should be a sufficient reason for referral of a child or young person, even when the primary healthcare professional considers that the symptoms are most likely to have a benign cause. [1.14.4]

Persistent back pain in a child or young person can be a symptom of cancer and is indication for an examination, investigation with a full blood count and blood film, and consideration of referral. [1.14.5]

There are associations between Down's syndrome and leukaemia, between neurofibromatosis and CNS tumours, and between other rare syndromes and some cancers. The primary healthcare professional should be alert to the potential significance of unexplained symptoms in children or young people with such syndromes. [1.14.6]

The primary healthcare professional should convey information to the parents and child/young person about the reason for referral and which service the child/young person is being referred to so that they know what to do and what will happen next. [1.14.7]

The primary healthcare professional should establish good communication with the parents and child/young person in order to develop the supportive relationship that will be required during the further management if the child/young person is found to have cancer. [1.14.8]

Leukaemia usually presents with a relatively short history of weeks rather than months. The presence of one or more of the following symptoms and signs requires investigation with full blood count and blood film:

- pallor
- fatigue
- unexplained irritability
- unexplained fever
- persistent or recurrent upper respiratory tract infections
- generalised lymphadenopathy
- persistent or unexplained bone pain

48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015] [R119]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. **[new 2015] [R120]**

Take into account the insight and knowledge of parents and carers when considering urgent referral for a child or young person. **[new 2015] [R121]**

Consider referral for children if their parent or carer has persistent concern or anxiety about the child's symptoms, even if the symptoms are most likely to have a benign cause. **[new 2015] [R122]**

Replaced by:
[R106]

Offer a very urgent full blood count (within 48 hours) to assess for leukaemia in children and young adults with any of the following symptoms:

- pallor
- persistent fatigue
- unexplained fever
- unexplained persistent infection
- generalised lymphadenopathy
- persistent or unexplained bone pain
- unexplained bruising
- unexplained bleeding. **[new 2015] [R106]**

- unexplained bruising.

If the blood film or full blood count indicates leukaemia then an urgent referral should be made. [1.14.9]

The presence of either of the following signs in a child or young person requires immediate referral:

- unexplained petechiae
- hepatosplenomegaly [1.14.10]

Lymphadenopathy is more frequently benign in younger children but urgent referral is advised if one or more of the following characteristics are present, particularly if there is no evidence of local infection:

- lymph nodes are non-tender, firm or hard
- lymph nodes are greater than 2 cm in size
- lymph nodes are progressively enlarging
- other features of general ill-health, fever or weight loss
- the axillary nodes are involved (in the absence of local infection or dermatitis)
- the supraclavicular nodes are involved. [1.14.11]

The presence of hepatosplenomegaly requires immediate referral. [1.14.12]

Shortness of breath is a symptom that can indicate chest involvement but may be confused with other conditions such as asthma.

Shortness of breath in association with the above signs (recommendation 1.14.11), particularly if not responding to bronchodilators, is an indication for urgent referral. [1.14.13]

A child or young person with a mediastinal or hilar mass on chest X-ray should be referred immediately. [1.14.14]

Persistent headache in a child or young person requires a neurological examination by the primary healthcare professional. An urgent referral should be made if the primary healthcare professional is unable to undertake an adequate examination. [1.14.15]

Headache and vomiting that cause early morning waking or occur on waking are classical signs of raised intracranial pressure, and an immediate referral should be

Replaced by:
[R111] – [R112]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for non-Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy or splenomegaly. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus or weight loss. **[new 2015]** **[R111]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for Hodgkin's lymphoma in people presenting with unexplained lymphadenopathy. When considering referral, take into account any associated symptoms, particularly fever, night sweats, shortness of breath, pruritus, weight loss or alcohol-induced lymph node pain. **[new 2015]** **[R112]**

Replaced by:
[R102] and [R104]

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for thyroid cancer in people with an unexplained thyroid lump. **[new 2015]** **[R102]**

Consider very urgent referral (for an appointment within 48 hours) for suspected brain or central nervous system cancer in children and young adults with newly abnormal cerebellar or other central neurological function. **[new 2015]** **[R104]**

made. [1.14.16]

The presence of any of the following neurological symptoms and signs should prompt urgent or immediate referral:

- new-onset seizures
- cranial nerve abnormalities
- visual disturbances
- gait abnormalities
- motor or sensory signs
- unexplained deteriorating school performance or developmental milestones
- unexplained behavioural and/or mood changes. [1.14.17]

A child or young person with a reduced level of consciousness requires emergency admission.[1.14.18]

In children aged younger than 2 years, any of the following symptoms may suggest a CNS tumour, and referral (a indicated below) is required.

- Immediate referral:
 - new-onset seizures
 - bulging fontanelle
 - extensor attacks
 - persistent vomiting.
- Urgent referral:
 - abnormal increase in head size
 - arrest or regression of motor development
 - altered behaviour
 - abnormal eye movements
 - lack of visual following
 - poor feeding/failure to thrive.
- Urgency contingent on other factors:
 - squint. [1.14.19]

Most children and young people with neuroblastoma have symptoms of metastatic disease which may be general in nature (malaise, pallor, bone pain, irritability, fever or respiratory symptoms), and may resemble those of acute leukaemia. The presence of any of the following symptoms and signs requires investigation with a full blood count:

- persistent or unexplained bone pain (and X-ray)
- pallor
- fatigue
- unexplained irritability
- unexplained fever
- persistent or recurrent upper

Replaced by:
[R117]

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]**
[R117]

- respiratory tract infections
- generalised lymphadenopathy
- unexplained bruising. [1.14.20]

Other symptoms which should raise concern about neuroblastoma and prompt urgent referral include:

- proptosis
- unexplained back pain
- leg weakness
- unexplained urinary retention. [1.14.21]

In children or young people with symptoms that could be explained by neuroblastoma, an abdominal examination (and/or urgent abdominal ultrasound) should be undertaken, and a chest X-ray and full blood count considered. If any mass is identified, an urgent referral should be made. [1.14.22] Infants aged younger than 1 year may have localised abdominal or thoracic masses, and in infants younger than 6 months of age, there may also be rapidly progressive intra-abdominal disease. Some babies may present with skin nodules. If any such mass is identified, an immediate referral should be made. [1.14.23]

Wilms' tumour most commonly presents with a painless abdominal mass. Persistent or progressive abdominal distension should prompt abdominal examination, and if a mass is found an immediate referral be made. If the child or young person is uncooperative and abdominal examination is not possible, referral for an urgent abdominal ultrasound should be considered. [1.14.24]

Haematuria in a child or young person, although a rarer presentation of a Wilms' tumour, merits urgent referral. [1.14.25]

A soft tissue sarcoma should be suspected and an urgent referral should be made for a child or young person with an unexplained mass at almost any site that has one or more of the following features. The mass is:

- deep to the fascia
- non-tender
- progressively enlarging
- associated with a regional

Replaced by:
[R119] – [R120]

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015] [R119]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. **[new 2015] [R120]**

Replaced by:
[R115] – [R116]

Consider an urgent direct access ultrasound scan (within 2 weeks) to assess for soft tissue sarcoma in people with an unexplained lump that is increasing in size. **[new 2015] [R115]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if they have ultrasound scan findings that are suggestive of soft

- lymph node that is enlarging
- greater than 2 cm in diameter. [1.14.26]

A soft tissue mass in an unusual location may give rise to misleading local and persistent unexplained symptoms and signs, and the possibility of sarcoma should be considered. These symptoms and signs include:

- head and neck sarcomas:
 - proptosis
 - persistent unexplained unilateral nasal obstruction with or without discharge and/or bleeding
 - aural polyps/discharge
- genitourinary tract:
 - urinary retention
 - scrotal swelling
 - bloodstained vaginal discharge.

[1.14.27]

Limbs are the most common site for bone tumours, especially around the knee in the case of osteosarcoma. Persistent localised bone pain and/or swelling requires an X-ray. If a bone tumour is suspected, an urgent referral should be made. [1.14.28]

History of an injury should not be assumed to exclude the possibility of a bone sarcoma. [1.14.29]

Rest pain, back pain and unexplained limp may all point to a bone tumour and require discussion with a paediatrician, referral or X-ray. [1.14.30]

In a child with a white pupillary reflex (leukocoria) noted by the parents, identified in photographs or found on examination, an urgent referral should be made. The primary healthcare professional should pay careful attention to the report by a parent of noticing an odd appearance in their child's eye. [1.14.31]

A child with a new squint or change in visual acuity should be referred. If cancer is suspected, referral should be urgent, but otherwise referral should be non-urgent. [1.14.32]

A family history of retinoblastoma should alert the primary healthcare professional to the possibility of retinoblastoma in a child who presents with visual problems. Offspring of a parent who has had retinoblastoma, or siblings of an

tissue sarcoma or if ultrasound findings are uncertain and clinical concern persists. **[new 2015] [R116]**

Replaced by:
[R113] – [R114]

Consider an urgent direct access X-ray (within 2 weeks) to assess for bone sarcoma in children and young adults with unexplained bone swelling or pain. **[new 2015] [R113]**

Consider a suspected cancer pathway referral (for an appointment within 2 weeks) for people if an X-ray suggests the possibility of bone sarcoma. **[new 2015] [R114]**

Replaced by:
[R118]

Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex. **[new 2015] [R118]**

affected child, should undergo screening soon after birth.

[1.14.33]

When cancer is suspected in children and young people, imaging is often required. This may be best performed by a paediatrician, following urgent or immediate referral by the primary healthcare professional. [1.14.34]
The presence of any of the following symptoms and signs requires investigation with full blood count:

- pallor
- fatigue
- irritability
- unexplained fever
- persistent or recurrent upper respiratory tract infections
- generalised lymphadenopathy
- persistent or unexplained bone pain (and X-ray)
- unexplained bruising. [1.14.35]

Replaced by:
[R117] – [R120]

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for neuroblastoma in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]** **[R117]**

Consider urgent referral (for an appointment within 2 weeks) for ophthalmological assessment for retinoblastoma in children with an absent red reflex. **[new 2015]** **[R118]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with a palpable abdominal mass or unexplained enlarged abdominal organ. **[new 2015]** **[R119]**

Consider very urgent referral (for an appointment within 48 hours) for specialist assessment for Wilm's tumour in children with unexplained visible haematuria. **[new 2015]** **[R120]**

Changes to recommendation wording for clarification only (no change to meaning)

Recommendation numbers in current guideline	Comment
All recommendations labelled [2005] .	Recommendations have been edited into the direct style (in line with current NICE style for recommendations in clinical guidelines) where possible. Yellow highlighting has not been applied to these changes.